



4% MALATHION DUST

ACCEPTED
Feb-5-1969
1384-220
96%

INGREDIENTS	
ACTIVE INGREDIENTS:	
Malathion*	4%
INERT INGREDIENTS	
Total	100%

*O-O-dimethyl dithiophosphate of diethyl mercaptosuccinate
USDA Reg. No. 1386-220

CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.
This product is poisonous to fish. Keep out of any body of water. Apply only in the manner and for the purposes stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.
DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE VEGETABLE CROPS

Begin applications before insects become numerous, or before, their damage is noticeable. Apply uniformly to cover upper and lower surfaces of all foliage. Repeat applications as necessary at 7 to 10-day intervals.
For the insects on the crops listed below, apply 35 to 40 pounds of dust (1.4 to 1.6 lb. malathion) per acre (equivalent to 4 ounces per 250 sq. ft. of crop area):
BROCCOLI, CABBAGE, CAULIFLOWER, KALE, TURNIP, MUSTARD—Aphids, imported cabbage worms, and diamond-back moth larvae.
BEANS—Mexican bean beetle, leafhoppers, spider mites.
CUCUMBER, SQUASH, MELON—Aphids, spider mites.
EGGPLANT—Aphids, spider mites.
LETTUCE—Aphids, spider mites, leaf hoppers.
ONIONS—Thrips.
POTATOES—Aphids.
TOMATOES—Spider mites, aphids.
For the insects on the crops listed below, apply 30 pounds of dust (1.20 lb. malathion) per acre (equivalent to 3 ounces per 250 sq. ft. of crop area):
PEAS—Pea aphid. PEPPERS—Aphids. SPINACH—Aphids.
DO NOT APPLY TO BROCCOLI, BEANS, SQUASH, EGGPLANT, ONIONS (GREEN), PEAS, PEPPERS, TOMATOES, OR TURNIP WITHIN 3 DAYS OF HARVEST.
DO NOT APPLY TO CABBAGE, CAULIFLOWER, KALE, MUSTARD, OR SPINACH WITHIN 3 DAYS OF HARVEST.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish. Keep out of any body of water. Apply only in the manner and for the purposes stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

VEGETABLE CROPS

Begin applications before insects become numerous, or before, their damage is noticeable. Apply uniformly to cover upper and lower surfaces of all foliage. Repeat applications as necessary at 7 to 10-day intervals.

For the insects on the crops listed below, apply 35 to 40 pounds of dust (1.4 to 1.6 lb. malathion) per acre (equivalent to 4 ounces per 250 sq. ft. of crop area):

BROCCOLI, CABBAGE, CAULIFLOWER, KALE, TURNIP, MUSTARD—Aphids, imported cabbage worms, and diamond-back moth larvae.

BEANS—Mexican bean beetle, leafhoppers, spider mites.

CUCUMBER, SQUASH, MELON—Aphids, spider mites.

EGGPLANT—Aphids, spider mites.

LETTUCE—Aphids, spider mites, leaf hoppers.

ONIONS—Thrips.

POTATOES—Aphids.

TOMATOES—Spider mites, aphids.

For the insects on the crops listed below, apply 30 pounds of dust (1.20 lb. malathion) per acre (equivalent to 3 ounces per 250 sq. ft. of crop area):

PEAS—Pea aphid. PEPPERS—Aphids. SPINACH—Aphids.

DO NOT APPLY TO BROCCOLI, BEANS, SQUASH, EGGPLANT, ONIONS (GREEN), PEAS, PEPPERS, TOMATOES, OR TURNIP WITHIN 3 DAYS OF HARVEST.

DO NOT APPLY TO CABBAGE, CAULIFLOWER, KALE, MUSTARD, OR SPINACH WITHIN 7 DAYS OF HARVEST.

DO NOT USE ON LEAF LETTUCE WITHIN 14 DAYS OF HARVEST AND HEAD LETTUCE WITHIN 7 DAYS OF HARVEST.

DO NOT APPLY TO CUCUMBER AND MELON WITHIN 1 DAY OF HARVEST.

MUSHROOMS

MUSHROOMS—To control phorid and sciarid flies (adults) and to aid in the control of pigmy mites in producing mushroom houses, apply this dust, following picking, uniformly at the rate of 1 3/4 pound per 2,000 square feet of bed surface (equiv. to 1.52 lb. malathion per acre), or 2 pounds per 60-foot single house.

DO NOT APPLY TO MUSHROOMS WITHIN 1 DAY OF HARVEST.

ORNAMENTAL PLANTS

Ornamental plants, roses, shrubs and some ornamental trees may be affected by aphids and spider mites which may be controlled by two or more thorough uniform full coverage applications of this dust. Apply to infested plants when insects first appear and make repeated applications as needed to protect the plants during the growing season. Do not use on Boston, Maidenhair, or Pteris ferns. May cause injury to Crassula.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

NET WEIGHT

50 POUNDS

FC-871-A1268

PROD 1039

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO