

GREENHOUSES - Use for control of two-spotted mites, mealybugs, and white fly on azaleas, camellias, gardenias, and other hardy plants. Use 1 1/2 to 2 tablespoons per gallon of water and apply spray at 7 a.m. to branches and foliage. Do not apply spray more than 60 days apart and do not make more than 3 applications per season.

TO MIX: For spray tank about one-third full with water, pour in the required amount of UNIC 3 Spray Oil, stir or agitate thoroughly and then fill remainder of tank with water. Some agitation is required to maintain the emulsion while spraying.

The dilution rate of 5 tablespoons per gallon is equivalent to 1 1/2 pint per 3 gallons or 2 gallons per 100 gallons of water.


CAUTION

Always use UNIC 3 Spray Oil in accordance with spray instructions. Do not apply to plants when temperatures are likely to drop below freezing soon after application. Do not spray on plants when near freezing temperatures or when the spray will freeze immediately before it dries. Weather conditions may vary and you may need to re-apply to some plants. Do not spray on plants when temperatures are likely to drop below freezing soon after application.

Use UNIC 3 Spray Oil in accordance with spray instructions. Do not apply to plants when temperatures are likely to drop below freezing soon after application. Do not spray on plants when near freezing temperatures or when the spray will freeze immediately before it dries. Weather conditions may vary and you may need to re-apply to some plants. Do not spray on plants when temperatures are likely to drop below freezing soon after application.

NOTICE

Use UNIC 3 Spray Oil in accordance with spray instructions. Do not apply to plants when temperatures are likely to drop below freezing soon after application. Do not spray on plants when near freezing temperatures or when the spray will freeze immediately before it dries. Weather conditions may vary and you may need to re-apply to some plants. Do not spray on plants when temperatures are likely to drop below freezing soon after application.



For Dormant and Foliage Applications on Fruit Trees, Deciduous Shade Trees, Hardy Ornamental Shrubs, and Plant Polish for Hardy House Plants.

INGREDIENTS

ACTIVE INGREDIENTS:		
Refined petroleum distillate		97%
NEEFT INGREDIENTS:		3%
Total Unsulfonated residue	92% minimum	100%
Class or grade	Unclassified	
Viscosity at 70°F	60-80	

CAUTION

KEEP OUT OF REACH OF CHILDREN.

See Remainder of CAUTION Statement on Side Panel.

EPA Reg. No. 1386-246

Net Volume 1 U.S. Quart Liquid

REG. 196-15-92

MANUFACTURED BY

PROD. 2126

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

FRUIT T...
 nsects and e
 worm on app
 of water and
 on bright, cl
 just before g

During t
 water and a
 two-spotted
 before they
 at 1-3 appli

SHADE T...
 period to co
 trees and har
 spray throug
 dry when th

Use UNIC 3
 on plants w
 temperatures
 near freezing
 when the spr

HARDY H...
 Use UNIC 3



SPRAY OIL

UNICO Spray Oil

DIRECTIONS FOR USE

FRUIT TREES During the dormant or delayed-dormant period, to control scale insects and eggs of European red mite, clover mite, apple rust mite, and fall canker-worm on apples, pears, cherries, peaches, and pums, use 5 tablespoonfuls per gallon of water and apply spray thoroughly to twigs, branches, and trunk early in the spring, on bright, clear days when the temperature is above 45° F. during the time from just before growth starts until the young leaves are not more than 1/2 inch in length.

During the summer when foliage is present, use 5 tablespoonfuls per gallon of water and apply spray thoroughly to foliage and branches to control scale insects, two-spotted mite, red bug, and thrips. Apply when insects first appear, preferably before they cause noticeable damage, and repeat at 30-day intervals, if necessary. (Do 3 applications if necessary) during the growing season.

SHADE TREES AND ORNAMENTAL SHRUBS During the dormant or delayed-dormant period, to control scale insects and eggs of European red mite on deciduous shade trees and hardy ornamental shrubs, use 5 tablespoonfuls per gallon of water and apply spray thoroughly to twigs, branches, and trunk early in the spring, on bright, clear days when the temperature is above 45° F., just before growth starts.

During the summer when foliage is present, use 5 tablespoonfuls per gallon of water and apply spray thoroughly to twigs, branches, and foliage to control scale insects, two-spotted mite, thrips, and two-spotted mites. Apply when insects first appear, preferably before they cause noticeable damage, and repeat at 30-day intervals if necessary. (Do 3 applications if necessary) during the growing season.

HARDY HOUSE PLANTS Apply general purpose and plant polish for hardy house plants, use 2 tablespoonfuls per gallon of water and apply spray lightly to foliage.

For Dormant and Foliage Applications on Fruit Trees, Deciduous Shade Trees, Hardy Ornamental Shrubs, and Plant Polish for Hardy House Plants.

INGREDIENTS

ACTIVE INGREDIENTS:		97%
Retined petroleum distillate		
INERT INGREDIENTS:		3%
Total		100%
Unulfonated residue	92% minimum	
Class or grade	Unclassified	
Viscosity at 100°F	60 Seconds	

CAUTION

KEEP OUT OF REACH OF CHILDREN.

See Remainder of CAUTION Statement on Side Panel.
EPA Reg No 1386-246

Net Volume 1 U.S. Quart Liquid

H&G-59C-D-72 MANUFACTURED BY PROD 2126

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601