



# 10% CHLORDANE GRANULAR

## CAUTION

Keep Out of Reach of Children

### INGREDIENTS

ACTIVE INGREDIENTS:	
Technical Chlordane*	10%
INERT INGREDIENTS	90%
Total	100%

\*Equivalent to 6% octachloro-4,7-methanotetrahydroindane and 4% related compounds

E.P.A. Reg. No. 1386-280

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust or vapor. In case of contact with skin, wash with soap and water. Do not contaminate feed and foodstuffs.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply in any manner not specified in this label. Do not apply when weather conditions favor drift of dust from area treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label. Clean application equipment and dispose of wastes, and unused chemical in a pit on non-crop land located well away from water supplies.

**DESTROY EMPTY CONTAINERS:** Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

### DIRECTIONS FOR USE

#### PREPLANTING SOIL APPLICATIONS — DIRECT APPLICATION TO SOIL BROADCAST TREATMENTS

Apply as a uniform broadcast treatment of the surface of the soil (previously prepared for planting) at the rate per acre indicated, promptly follow with cultivation to mix the granules into the soil to a depth of 6 to 8 inches, preferably 7 to 14 days prior to planting or transplanting of the crops shown below. Properly applied broadcast preplanting soil treatments may be effective for more than one year; repeat applications as new infestations develop. Use the amount shown after each crop to control the insects named.

**CORN** — To control corn rootworms and seed corn maggot, 30 to 40 pounds per acre. To control wireworms and white grubs, 50 to 80 pounds per acre.

**TOBACCO** — To control wireworms, white grubs, and green June beetle larvae, 50 to 60 pounds per acre.

**POTATOES** — To control wireworms and white grubs, 50 to 80 pounds per acre.

**STRAWBERRIES** — To control wireworms, 50 to 80 pounds per acre; to control white grubs, 100 pounds per acre.

#### IN-ROW TREATMENTS

Apply uniformly in a band 10 to 12 inches wide centered over the row, promptly work the insecticide into the top 6 inches of soil, preferably 7 days prior to planting and transplanting of the crops named below. In-row treatments should be repeated annually. Use the amount shown after each crop to control the insect named.

**CORN** — To control corn rootworms, and seed corn maggot, 20 pounds per acre or 1 ounce per 136 square feet or 1 ounce per 136 feet of 12 inch band. To control wireworms and white grubs, 25 to 40 pounds per acre or 1 ounce per 82 square feet or 1 ounce per 82 feet of 12 inch band. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**TOBACCO** — To control wireworms and white grubs, 25 to 40 pounds per acre or 1 ounce per 82 square feet or 1 ounce per 82 feet of 12 inch band.

**POTATOES** — To control wireworms and white grubs, 25 to 40 pounds per acre or 1 ounce per 82 square feet or 1 ounce per 82 square feet of 12 inch band.

