

INGREDIENTS:

ACTIVE INGREDIENTS:

Technical Chlordane* 5.0% INERT INGREDIENTS 95.0% Total 100.0% *Equivalent to 3.0% octachloro-4,7-methanotetrahydroindane and 2* related compounds EPA REG. NO. 1386-287

CAUTION Keep Out Of Reach Of Children

Harmful if swallowed. Avoid breathing dust and vapor. Avoid contamination of feed and foodstuffs. Highly toxic to fish and wildlife. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on this label. This product is toxic to fish and wildlife. Birds feeding on the treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay

away from smoke or fumes.

DIRECTIONS FOR USE

PREPLANTING SOIL APPLICATIONS

CUTWORMS — Apply 25 to 30 lb. per acre (3 ounces per 250 sq. ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper, or tobacco transplants. If desired, place 2 to 3 teaspoonfuls in a ring around each transplant but not in direct contact with the plant.

WIREWORMS — Apply 75 to 100 lb. per acre (1 to 11/4 lb. per 500 sq. ft.) uniformly over surface of prepared land, disk lightly into top 3 or 4 inches of soil ahead of planting of potatoes, corn, sweet corn, tobacco, or strawberries.

CABBAGE ROOT MAGGOT — Apply 75 to 100 lb. per acre (1 to 11/4 lb. per 500 sq. ft.). Work into top 3 inches of soil just before setting of cabbage, broccoli, or cauliflower transplants.

SEED CORN MAGGOT - Apply 40 lb. per acre (4 ounces per 250 sq. ft.). Work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS — Uniformly apply 150 to 200 lb. per acre (3 ounces per 250 sq. ft.) of prepared soil and disk into top 4 to 6 inches of soil just before setting of strawberry or ornamental transplants.

Repeat preplanting soil treatments annually if needed. In some situations, properly applied broadcast soil treatments may be effective for more than one year. LAWNS AND ORNAMENTAL TURF

ANTS, CHIGGERS, CRICKETS, AND EARWIGS — Apply 25 to 30 lb. per acre (3 ounces per

250 sq. ft. of area) directly into the grass. Repeat as necessary. CUTWORMS — Apply at rate of 30 to 40 lbs. per acre.

SOD WEBWORMS — Apply at rate of 40 lbs. per acre. For imported fire ant, break open hard surface of mound and pour in 2 to 4 cupfuls per mound; sprinkle surrounding soil as for other ants. If ants survive, repeat in two weeks.

INGREDIENTS:

ACTIVE INGREDIENTS: Technical Chlordane* INERT INGREDIENTS

Total *Equivalent to 3.0% octachloro-4,7-methanotetrahydroindane and 2% related compounds EPA REG. NO. 1386-287

CAUTION Keep Out Of Reach Of Children

Harmful if swallowed. Avoid breathing dust and vapor. Avoid contamination of feed and foodstuffs. Highly toxic to fish and wildlife. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on this label. This product is toxic to fish and wildlife. Birds feeding on the treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

PREPLANTING SOIL APPLICATIONS

CUTWORMS — Apply 25 to 30 lb. per acre (3 ounces per 250 sq. ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper, or tobacco transplants. If desired, place 2 to 3 teaspoonfuls in a ring around each transplant but not in direct contact with the plant.

WIREWORMS — Apply 75 to 100 lb. per acre (1 to 1¼ lb. per 500 sq. ft.) uniformly over surface of prepared land, disk lightly into top 3 or 4 inches of soil ahead of planting of potatoes, corn, sweet corn, tobacco, or strawberries.

CABBAGE ROOT MAGGOT — Apply 75 to 100 lb. per acre (1 to $1\frac{1}{4}$ lb. per 500 sq. ft.). Work into top 3 inches of soil just before setting of cabbage, broccoli, or cauliflower transplants.

SEED CORN MAGGOT — Apply 40 lb. per acre (4 ounces per 250 sq. ft.). Work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS — Unifomly apply 150 to 200 lb. per acre (3 ounces per 250 sq. ft.) of prepared soil and disk into top 4 to 6 inches of soil just before setting of strawberry or ornamental transplants.

Repeat preplanting soil treatments annually if needed. In some situations, properly applied broadcast soil treatments may be effective for more than one year.

LAWNS AND ORNAMENTAL TURF

ANTS, CHIGGERS, CRICKETS, AND EARWIGS — Apply 25 to 30 lb. per acre (3 ounces per 250 sq. ft. of area) directly into the grass. Repeat as necessary. CUTWORMS — Apply at rate of 30 to 40 lbs. per acre.

SOD WEBWORMS — Apply at rate of 40 lbs. per acre.

For imported fire ant, break open hard surface of mound and pour in 2 to 4 cupfuls per mound; sprinkle surrounding soil as for other ants. If ants survive, repeat in two weeks. WHITE GRUBS AND JAPANESE BEETLE GRUBS — Apply 150 lb. per acre (1 lb. per 250

sq. ft. of area) in early spring or late fall.

Keep children and pets off treated areas until the insecticide has been thoroughly washed into the soil, either by rainfall or thorough sprinkling, and the grass has completely dried. Do not feed grass from treated areas to poultry, dairy animals, or animals being finished for slaughter. Do not pasture such animals on treated areas. Birds feeding over treated areas may be harmed.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

NET WEIGHT

FC-868-A4-72

DISTRIBUTED BY UNITED CO-OPERATIVES, INC. ALLIANCE, OHIO 44601







should be injected into the soil beneath the floor. Avoid drilling into plumbing or electric conduits. The emulsion should be applied at the rate of at least 4 gailons per 10 linear feet of wall.

(3) Treat voids in hollow-block foundations at the rate of 1 gallon per 5 linear feet of wall so that the emulsion will reach the footing. Do this by drilling or probing. Take care to avoid drilling into plumbing or electric conduits.

Slab-on-ground Construction:

(1) Infestations in this type of construction are difficult to control. One method consists of drilling holes about a foot apart through the concrete slab, adjacent to all cracks and expansion joints, and injecting the chemical into the soil beneath the slab. Avoid drilling into plumbing and electric conduits. Another method is to drill through the foundation walls from the outside and force the chemical just beneath the slab along the inside of the foundation and along all the cracks and expansion joints. The emulsion should be applied at the rate of at least 4 gallons per 10 linear feet of foundation or expansion joints.

(2) Dig a trench 1 foot in depth, but not below the top of the footing, alon_n the outside of the foundation walls. Apply the emulsion at the rate of 4 gallons per 10 linear feet of trench. The chemical should be mixed with the soil as it is being replaced in the trench.

(3) Treat voids in hollow block foundations at the rate of one gallon of emulsion per 5 linear feet of wall so that the emulsion will reach the footing. Do this by drilling or probing. Avoid drilling into plumbing or electric conduits.

CAUTION

Harmful if swallowed! Contact with skin causes toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

DO NOT USE, SPILL, POUR OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, r ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

UNICO TERMITE KILL DIRECTIONS FOR USE

To control termites, remove all wood and other cellulose material in contact with the soil. Prepare a water emulsion containing 1 percent chlordane by mixing 1 pint of this concentrate in 6¼ gallons of water (1 gallon in 50 gallons), and apply as follows.

Preconstruction Treatment

Slab-on-ground Construction:

(1) Apply an overall treatment under entire surface of floor slab. Apply at the rate of 1 gallon per 10 square feet, except that if fill under slab is gravel or other coarse absorbent material, apply at the rate of 11/2 gallons per 10 square feet.

(2) Under slab-on-ground porch floors and entrance platforms, apply an overall treatment at the rate of 1 gallon per 10 savare feet.

(3) Along both sides of foundation wall, along interior foundation walls, and around plumbing dig a narrow trench to a depth of 1 foot, but not below the top of the footing. Apply at the rate of 2 gallons per 5 linear feet of trench. The chemical should be mixed with the soil as it is being replaced in the trench. (4) Treat all voids in hollow masonry units of the foundation at the rate of at least 1 gallon per 5 linear feet of wall. Apply the emulsion so as to reach the footing.

Buildings with Crawl Spaces:

(1) Dig a narrow trench to the top of the footing along the inside of foundation walls, around piers, sewer pipes and conduits. Apply 2 gallons of emulsion per 5 linear feet of trench. The chemical should be mixed with the soil as it is being replaced in the trench.

(2) Dig a narrow trench to the top of the footing along the ourside of the foundation wall. Apply 2 gallons of emulsion per 5 linear feet of trench per each foot of depth. A trench 3 feet deep would require 6 gallons per 5 linear feet. The chemical should be mixed with the will as it is being replaced in the trench. (3) Under attached porches, entrance platforms, utility entrances, and similar situations where slab or fill is at the same grade level apply 1 gallon per 10 square feet of soil surface. (4) Treat all voids in hollow masorry units of the foundation at the rate of at least 1 gallon per 5 linear feet of wall. Apply the emulsion so as to reach the footing.

Buildings with Basements:

(1) Apply an overall treatment under the basement floorings, as well as under attached porches, entrance platforms, utility entrances, and similar situations where slab fill is at the grade level. Apply at the rate of 1 gallon per 10 square feet, except that if fill under slab is of washed gravel, cinders or similar coarse material, increase the dosage by at least one-half. Where crawl spaces exist, treat as described in part (2) below. (2) Dig a narrow trench to the top of the footing along the inside of foundation walls, around piers, sewer pipes and conduits. Apply 2 gallons of emulsion per 5 linear feet of trench. The chemical should be mixed with the soil as it is being replaced in the trench.

(3) Along the outside of the foundation walls, dig a narrow trench, such trench to be dug no deeper than the top of the footings. If the trench is less than 15 inches in depth to the top of the footings, apply 1 gallon per 5 linear feet. Replace the soil and apply another 1 gallon per 5 linear feet to the back fill. Cover the back fill with a thin layer of soil. If the trench is more than 15 inches in depth to the top of the footings, apply 2 gallons per 5 linear feet. Replace the soil and apply another 2 gallons per 5 linear feet to the back fill. Cover the back fill with a thin layer of soil. A trench 30 inches deep is a maximum depth required alongside foundations where the top of the footings is greater than 30 inches deep. In lieu of trenching to a 30" depth, make the trench 12 to 15" deep and rod to footing, spacing the holes about 1 foot apart.

(4) Treat all voids in hollow masoury units of the foundation at the rate of 1 gallon per 5 linear feet of wall. Apply the emulsion so as to reach the footing.

Control in Existing Buildings

Buildings Having Crawl Spaces

(1) Dig a trench adjacent to and around all piers and pipes and along both sides of the foundation walls. Dig the trench to, but not below the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet. (2) Treat voids in hollow-block mast nry foundations at the rate of 1 gallon per 5 linear feet of wall. Apply so that the emulsion will reach the footing. 'f this is done by drilling or rod holes avoid going into plumbing or electrical conduits. **Buildings Having Basements**

(1) Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench isrefilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet. (2) It may also be necessary to treat critical areas only under the basement flooring such as around sewer pipes, conduits and piers and along the inside of the foundation walls and interior walls. One method consists of drilling holes about a foot apart through the concrete floor adjacent to the areas requiring treatment. The chemical emulsion then