

UNICO

SPINACH — Flea beetles, leafhoppers, downy mildew.
DO NOT APPLY WITHIN 14 DAYS OF HARVEST.

RADISHES AND TURNIPS — Flea beetles, alternaria and
downy mildew DO NOT APPLY WITHIN 3 DAYS OF HAR-
VEST. 14 DAYS IF TOPS ARE TO BE USED AS FOOD OR
FEED.

TOMATOES — Aphids, Colorado potato beetle, flea
beetle, leafhoppers, tomato fruit worm, tarnished plant
bug, anthracnose, early blight, late blight and Septoria
leaf spot DO NOT APPLY WITHIN 1 DAY OF HARVEST.

STRAWBERRIES — Spittlebug, strawberry leaf roller,
leafhoppers, aphid, strawberry weevil, Botrytis fruit rot,
leaf spot. Apply 40 pounds per acre. Sunrise and Early
Dawn varieties may be injured by Sevin.

Apply 1 when new growth starts in the spring, (2)
at subsequent 7 to 10 day intervals as needed. DO NOT
APPLY WITHIN 7 DAYS OF HARVEST.

CAUTION

HARMFUL IF INHALED OR SWALLOWED may cause irritation of skin, eyes, nose, throat or lungs. Avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after work; change clothing.

This product is poisonous to fish or wildlife. Keep out of
the body or water. Do not apply to crops or areas other
than those stated in this label and not when weather con-
ditions favor drift of dust from area treated. Clean
application equipment and dispose of wastes and unused
chemical material on non-crop land well away from water
sources. Avoid contamination of feed and foodstuffs.

DESTROY EMPTY CONTAINERS. Do not reuse this con-
tainer for any purpose. Destroy container in compliance
with local regulations. If burned stay away from smoke.

NETS — This product is highly toxic to bees exposed to
direct spray or to flowers of crops. Protection information
is available from the Cooperative Agricultural Ex-
periment Station.

NOTE FOR PETS — Can cause moderate to severe
irritation to animals. Atropine is an antidote.

NOTICE

Use only for the purposes and in compliance with the
instructions given on the reverse side of this label.



Garden Dust

INGREDIENTS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N methylcarbamate)	3.0%
Zineb (zinc ethylene bisdithiocarbamate)	3.2%
Ziram (zinc dimethyl dithiocarbamate)	3.8%
Rotenone	1.0%
Other cube resins	1.6%

INERT INGREDIENTS

Total 100.0%

3.0%
3.2%
3.8%
1.0%
1.6%
87.4%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

USDA Reg. No. 1386 316

NET WEIGHT

H&G 110

1 POUND

Prod. 2143

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

UNICO GARDEN DUST

DIRECTIONS FOR USE

Use on crops listed below to control the insects and plant diseases indicated. Begin applications before diseases are apparent and before insects become numerous which is usually when the seedlings appear above ground or young plants are transplanted to the garden. Repeat applications at 5 to 7 day intervals and after heavy rains to maintain protection during growing season. Cover foliage both upper and lower surfaces evenly and thoroughly with a light film of dust when air is calm. Apply at the rate of 35-40 pounds per acre 1 lb per 1,000 sq ft or 1 lb per 1,000 linear feet of row. Excessive application may cause scalding of tender foliage.

CROPS, INSECTS, AND PLANT DISEASES

BEANS — Aphids, anthracnose, white mold, Mexican bean beetle, bean leaf beetle, leafhoppers DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

BROCCOLI, CABBAGE, CALIFLOWER AND KALE — Imported cabbage worm, diamond back moth larvae, Har-
quin bug, and downy mildew DO NOT APPLY WITHIN
7 DAYS OF HARVEST.

CARROTS — Flea beetles and leafhoppers, early and late
blight DO NOT APPLY WITHIN 7 DAYS OF HARVEST IF
TOPS ARE TO BE USED AS FOOD OR FEED. OTHERWISE
DO NOT APPLY WITHIN 1 DAY OF HARVEST.

CUCUMBER, MELONS AND SQUASH — Aphids, leaf-
hoppers, melon worm, pickle worm, angular leaf spot,
downy mildew, crab and anthracnose DO NOT APPLY
WITHIN 1 DAY OF HARVEST.

Eggplant — Aphids, flea beetles, tomato hornworm,
angular leaf spot, early and late blight DO NOT APPLY
WITHIN 1 DAY OF HARVEST.

GARDEN BEETS — Flea beetles DO NOT APPLY
WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED
AS FOOD OR FEED. OTHERWISE DO NOT APPLY WITHIN
7 DAYS OF HARVEST.

LETTUCE — Downy mildew, flea beetles, leafhoppers
DO NOT APPLY WITHIN 14 DAYS OF HARVEST.

PEAS — Leafhoppers, aphids DO NOT APPLY WITHIN
10 DAYS OF HARVEST.

PEPPERS — Flea beetles, imported cabbage moths
and downy mildew DO NOT APPLY WITHIN 10 DAYS OF HARVEST.