



9-26-72
1386-325

WIREWORMS—Apply 35 to 50 lb. per acre (1/2 to 2/3 lb. per 500 sq. ft.) uniformly over surface of prepared land, work into top 3 or 4 inches of soil prior to planting of corn, sweet corn, tobacco, or strawberries. Does not include resistant wireworms or Southern potato wireworm.

SEED CORN MAGGOT — Apply 25 pounds per acre (5 oz. per 500 sq. ft.). Work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS—Uniformly apply 75 to 100 lb. per acre (1 lb. per 500 sq. ft.) of prepared soil and work into top 4 to 6 inches of soil just before setting of strawberry or ornamental transplants. Do not apply to strawberries after transplanting.

CAUTION

Avoid breathing dust or vapor. Avoid contamination of feed and foodstuffs.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

10% CHLORDANE Dust

For Application to Lawns and Soil

INGREDIENTS

ACTIVE INGREDIENTS:		
Technical Chlordane*	10.0%
INERT INGREDIENTS	90.0%
		100.0%
Total	100.0%

*Equivalent to 6% octachloro-4,7-methanctetrahydroindane and 4% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statements on Side Panel

EPA Reg. No. 1386-325

NET WEIGHT

H&G-102H-6-72

1 POUND

PROD. 2134

DISTRIBUTED BY

UNICO 10% Chlordane Dust

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF

ANTS, CHIGGERS, SOD WEBWORMS, CRICKETS, CUTWORMS, AND EARWIGS—Apply 30 pounds per acre (5 1/2 oz. per 500 sq. ft. of area) directly into the grass. Repeat as necessary.

For imported fire ant in lawns and ornamental turf, break open hard surface of mound and pour in 1 to 2 cupfuls per mound; sprinkle surrounding soil as for other ants. If ants survive, repeat in 2 weeks.

WHITE GRUBS AND JAPANESE BEETLE GRUBS—Apply 75 to 100 lb. per acre (1 lb. per 500 sq. ft. of area) in early spring or late fall. Repeat annually if necessary.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by adequate rainfall or thorough sprinkling, and the grass has completely dried. DO NOT feed or graze forage from treated areas to livestock or poultry. Birds feeding over treated areas may be harmed.

PREPLANTING SOIL APPLICATIONS

CUTWORMS — Apply 25 pounds per acre (5 oz. per 500 sq. ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper, or tobacco transplants. If desired, place 1 to 1 1/2 teaspoonfuls in a ring around each transplant but not in direct contact with plant.

UNITED CO-OPERATIVES INC.,
ALLIANCE, OHIO 44601