

UNICO

UNICO Methoxychlor Emulsifiable Concentrate

DIRECTIONS FOR USE

This product is a systemic insecticide for control of sucking insects on a wide range of ornamental plants, shrubs, trees, and lawns. For complete directions, see the following paragraphs.

ALFALFA, CLOVER, AND GRASS - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. For control of alfalfa cutworm, apply to alfalfa and cut at 100-150 lbs. per acre. For grasshopper control, apply to alfalfa at 100-150 lbs. per acre. For clover leaf weevil control, apply to clover at 100-150 lbs. per acre. Apply in the evening and repeat in 7 days.


DO NOT APPLY TO ALFALFA OR GRASS TO BE HARVESTED WITHIN 7 DAYS OF HARVEST OR GRAZING.

SOYBEANS, COWPEAS, AND BEANS - Control soybean and cowpea weevils, and bean beetles. Apply 1.0 to 2.0 fl. oz. per acre. For soybean and cowpea weevil control, apply to soybeans and cowpeas at 100-150 lbs. per acre. For bean beetle control, apply to beans at 100-150 lbs. per acre. Apply in the evening and repeat in 7 days.

DO NOT APPLY TO SOYBEANS, COWPEAS, OR BEANS WITHIN 7 DAYS OF HARVEST OR GRAZING.

VEGETABLE CROPS - Control alfalfa cutworm, clover leaf weevil, grasshopper, and cowpea weevil. Apply 1.0 to 2.0 fl. oz. per acre. For alfalfa cutworm control, apply to alfalfa and cut at 100-150 lbs. per acre. For clover leaf weevil control, apply to clover at 100-150 lbs. per acre. For grasshopper control, apply to alfalfa at 100-150 lbs. per acre. For cowpea weevil control, apply to cowpeas at 100-150 lbs. per acre. Apply in the evening and repeat in 7 days.

- ASPARAGUS - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- BEANS - Control bean beetle. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- PEAS - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- CUCUMBERS, CANTALOUPES, PUMPKINS, SUMMER AND WINTER SQUASH - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- EGGPLANTS - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- LETTUCE - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- POTATOES - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.
- TOMATOES - Control alfalfa cutworm, clover leaf weevil, and grasshopper. Apply 1.0 to 2.0 fl. oz. per acre. Apply in the evening and repeat in 7 days.



INGREDIENTS

ACTIVE INGREDIENT:
Methoxychlor (95% min)
Aromatic petroleum hydrocarbon solvent
INERT INGREDIENTS

Total

*Equivalent to 20.9% 2,2-bis(p-methoxyphenyl)propane and 2.9% other isomers and related compounds.

CONTAINS 2.5% TECHNICAL METHOXYCHLOR

WARNING

KEEP OUT OF REACH OF CHILDREN

See Remainder of WARNING Statement on Label

EPA Reg. No. 1386-0
EPA Est. No. 1386-0001

Net Volume

5 U.S. GALS.

ACCEPTED
AUG 13 1966
U.S. DEPT. OF AGRICULTURE
EPA Reg. No. 1386-0
EPA Est. No. 1386-0001

TOPS ARE TO BE STORED IN A COOL PLACE, SUCH AS A REFRIGERATOR, WITHIN 24 HOURS OF HARVEST. DO NOT WASH OR TRIM TOPS UNTIL JUST BEFORE USE. TOPS ARE TO BE STORED IN A COOL PLACE, SUCH AS A REFRIGERATOR, WITHIN 24 HOURS OF HARVEST. DO NOT WASH OR TRIM TOPS UNTIL JUST BEFORE USE.

GRAIN STORAGE BINS

Grain storage bins should be inspected regularly for signs of infestation. If infestation is detected, the grain should be treated with an appropriate insecticide. The grain should be stored in a cool, dry place and should be protected from moisture. The grain should be stored in a cool, dry place and should be protected from moisture.

Grain storage bins should be inspected regularly for signs of infestation. If infestation is detected, the grain should be treated with an appropriate insecticide. The grain should be stored in a cool, dry place and should be protected from moisture.

Grain storage bins should be inspected regularly for signs of infestation. If infestation is detected, the grain should be treated with an appropriate insecticide. The grain should be stored in a cool, dry place and should be protected from moisture.

BEE, CATTLE, SHEEP, GOATS, AND HOGS

Bees, cattle, sheep, goats, and hogs should be handled with care. Bees should be kept in a secure location. Cattle, sheep, goats, and hogs should be kept in a secure location and should be protected from predators. The animals should be kept in a secure location and should be protected from predators.

Bees, cattle, sheep, goats, and hogs should be handled with care. Bees should be kept in a secure location. Cattle, sheep, goats, and hogs should be kept in a secure location and should be protected from predators.

WARNING

Warning: This product is highly flammable. Keep away from fire and heat. Do not use near open flames. Use in a well-ventilated area. Avoid contact with skin and eyes. If contact occurs, wash immediately with water. For more information, see the Safety Data Sheet (SDS).

Warning: This product is highly flammable. Keep away from fire and heat. Do not use near open flames. Use in a well-ventilated area. Avoid contact with skin and eyes. If contact occurs, wash immediately with water. For more information, see the Safety Data Sheet (SDS).

NOTICE

Notice: This product is highly flammable. Keep away from fire and heat. Do not use near open flames. Use in a well-ventilated area. Avoid contact with skin and eyes. If contact occurs, wash immediately with water. For more information, see the Safety Data Sheet (SDS).