

Methoxychlor
Emulsifiable Concentrate

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENTS:

Methoxychlor: 1,1,1-trichloro-2, 2-bis(p-methoxyphenyl) ethane*	25.0%
INERT INGREDIENTS**	75.0%
Total	100.0%

*Equivalent to 21.9% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloro-ethane and 3.1% other isomers and related compounds. Contains 2 pounds technical methoxychlor per gallon. **Contains Aromatic Petroleum Derivative Solvent.

Net Volume: 1 Gallon

EPA Reg. No. 1386-352
EPA Est. No. 1386-01-1

BEST AVAILABLE COPY



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN, wash with soap and water. Remove contaminated clothing and wash before reuse. **IF IN EYES**, flush eyes with plenty of water. Get medical attention. **IF INHALED**, remove victim to fresh air. Apply respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other wildlife. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is an emulsifiable insecticide to be mixed with water (forming an emulsion) for application as a spray to control certain insects on the following paragraphs.

BEEF CATTLE, SHEEP, GOATS AND HOGS—To control horn flies and lice, use 1 gallon mixed in 48 gallons of water (1 qt. in 12 gal.) to make a spray containing 0.5% methoxychlor. Apply as a spray only. Do not use in dipping vats. Apply the spray thoroughly to each animal, using 2 quarts per acre for large beef cattle and proportionally less on smaller animals, sheep, goats and hogs. Repeat treatment in 2 to 3 weeks, or as needed.

DO NOT APPLY TO DAIRY ANIMALS OR TO LACTATING GOATS.

GRAIN STORAGE BINS—To control accessible stages of certain insect pests of stored grain which remain in empty storage bins and other storage areas, such as cadelle, rice weevil, granary weevil, saw-toothed grain beetle, lesser grain borer, flat grain beetle, confused flour beetle and other beetles, use 1 gallon of this concentrate mix in 10 gallons of water (1 quart in 2½ gallons of water) and spray inside and outside surfaces of empty granaries and other storage facilities prior to filling with grain. Apply the spray at the rate of 1 gallon per 500 square feet to all surfaces on which insects rest or crawl, or where they rest or hide, using a coarse driving spray to penetrate into cracks and joints. Repeat as necessary.

DO NOT ADD GRAIN TO SPRAYED BINS OR OTHER STORAGE AREAS FOR AT LEAST 24 HOURS OR UNTIL THE SPRAYED SURFACES HAVE DRIED THOROUGHLY.

THOROUGHLY CLEAN BINS AND OTHER STORAGE AREAS ALWAYS PRIOR TO SPRAYING. Maintain good housekeeping and sanitation in storage areas as a distinct aid in preventing and reducing insect infestations.

CROP USES

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more frequent reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Warnings must inform workers of areas or fields that may not be entered without wearing specific protective clothing until sprays have dried and appropriate actions to take in case of accidental exposure (refer to Statement of Practical Treatment). When oral warnings are given, warnings shall be in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION. Area treated with Methoxychlor on (insert date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, follow these procedures: (insert information from Statement of Practical Treatment)."

ALFALFA, CLOVER AND GRASS—To control spittlebug, leafhoppers, clover leaf weevil and alfalfa caterpillar, use 2 quarts (1.0 lb. methoxychlor) per acre. To control alfalfa weevil (larvae), alfalfa webworm, Lygus bugs, and fall armyworm, use 3 quarts (1.5 lbs. methoxychlor) per acre. Apply in the amount of water required for uniform thorough spray coverage which is usually 15 to 20 gallons per acre for boom sprayers. Begin application when insects first appear, usually in the spring, and repeat as necessary.

DO NOT APPLY TO ALFALFA, CLOVER AND GRASS FORAGE WITHIN 7 DAYS OF HARVEST OR GRAZING.

SOYBEANS, COWPEAS AND PEANUTS—To control velvet bean caterpillar and soybean caterpillar, use 2 to 3 quarts (1.0 to 1.5 lb. methoxychlor) per acre. To control Mexican bean beetle, Japanese beetle (adults), pea weevil, cowpea curculio, alfalfa looper and celery looper, use 3½ quarts (1.75 lbs. methoxychlor) per acre. Apply in the amount of water required for uniform thorough spray coverage, which is usually 15 to 20 gallons per acre for boom sprayers. Begin applications when insects first appear, or as soon as feeding is noticed, and repeat as necessary.

DO NOT APPLY TO SOYBEANS, COWPEAS AND PEANUTS WITHIN 7 DAYS OF HARVEST OR GRAZING.

VEGETABLE CROPS—To control the insects on the crops listed below, use 3 to 3½ quarts (1.5 to 1.75 lbs. methoxychlor) per acre in the amount of water required for uniformly thorough spraying of one acre. Begin applications when insects first appear and repeat at 7 to 10 day intervals as necessary.

ASPARAGUS—Asparagus beetles.

BEANS—Mexican bean beetle, bean leaf beetle and leafhoppers.

BETTS—Blister beetles and flea beetles.

CUCUMBERS, CANTALOUPE, PUMPKINS, SUMMER AND WINTER SQUASH—Cucumber beetles.

EGGPLANTS—Flea beetles and leafhoppers.

LETTUCE—Leafhoppers.

POTATOES—Colorado potato beetle, leafhoppers and flea beetles.

TOMATOES—Blister beetles, flea beetles, fall armyworm, Japanese beetle (adults) and tomato fruitworm.

DO NOT APPLY TO CUCUMBERS, CANTALOUPE, EGGPLANTS, PUMPKINS, SUMMER SQUASH, WINTER SQUASH AND TOMATOES WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO ASPARAGUS WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO BEETS WITHIN 7 DAYS OF HARVEST (14 DAYS OF HARVEST). DO NOT APPLY TO LETTUCE WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO TOMATOES WITHIN 14 DAYS OF HARVEST.

STORAGE AND DISPOSAL



STORAGE: Do not store near heat or open flame. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures approved by State and local authorities.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so. This warranty is given with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material to which a claim is made.



BEST AVAILABLE COPY