

afternoon, as needed. Do not apply as a barn spray if animals which have received direct application within 8 hours remain in the barn.

DO NOT CONTAMINATE FEED, DRINKING WATER, MILK, AND MILKING EQUIPMENT.

### CAUTION

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. Avoid inhalation of vapor or spray mist. Avoid contact with skin, eyes, and clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate water, feed, or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

### NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

### USE THE UNICO 580U SPRAYER

THE UNICO 580U 3 QUART CONTINUOUS SPRAYER FEATURES AN ADJUSTABLE NOZZLE WHICH WILL DELIVER A FINE OR A COARSE SPRAY. UNICO PY-VONA PERFORMS IDEALLY IN THIS SPRAYER.



# PY-VONA<sup>®</sup> STOCK FLY SPRAY

Contains Synergized Pyrethrins and DDVP

ACTIVE INGREDIENTS	100%
Petroleum base oil	99.52%
2,2-Dichlorovinyl dimethyl phosphate	0.18%
Related compounds	0.02%
Technical piperonyl butoxide*	0.25%
Pyrethrins	0.03%
*Equivalent to 0.20% (butyl carbityl) (6-propyl piperonyl) ether and 0.05% related compounds	

### CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Required Precautionary Labeling.

USDA Reg. No. 1386-418

Net Volume 1 U.S. Quart Liquid

FC-749A-1-65

E-OD-3

MANUFACTURED BY

UNITED CO-OPERATIVES, INC.  
ALLIANCE, OHIO

## UNICO PY-VONA Stock Fly Spray

### DIRECTIONS FOR USE

UNICO PY-VONA Stock Fly Spray contains DDVP, piperonyl butoxide, and pyrethrins in a mineral seal oil base. It is a ready-to-use spray for application directly to dairy cattle, beef cattle, hogs, sheep, goats, and horses to control house flies, horn flies, stable flies, and mosquitoes, and to aid in control of face flies.

**DIRECT SPRAYING OF LIVESTOCK**—Apply as a finely atomized spray uniformly distributed over each animal at a rate of 1 to 2 fluid ounces per animal on dairy cattle, beef cattle, and horses, and at the rate of 1 fluid ounce per animal on hogs, sheep and goats. Reduce application rate by one-half for young animals. Use good spray equipment to spray each animal, preferably from the rear, directing the spray along both sides, including the belly, flanks, and withers, down each leg, over the back and around the poll. Hold nozzle of sprayer 2 or 3 feet away from animal, so as to deposit droplets on the hair. **DO NOT WET THE HIDE.** For milking dairy animals, apply the spray 20 to 30 minutes before beginning to milk, or after milking has been completed. **DO NOT SPRAY UDDERS OF LACTATING DAIRY ANIMALS. DO NOT APPLY IN EXCESS OF RATES STATED PER ANIMAL.** Repeat daily, as needed.

**BARN SPRAYING** To control house flies in dairy barns, milk rooms, loafing sheds, livestock barns, stables, and poultry houses, apply 2 fluid ounces per 1,000 cubic feet as a finely atomized spray, uniformly distributed throughout the area. To reduce air movement as much as possible, it is preferable to close windows and doors before spraying; leave them closed for 10 minutes after spraying and then ventilate the area. Repeat daily, morning and