TX3T W3ハ=柳緑

PY-VONA®

Stock Fly Spray

Contains Synergized Pyrethrins and DDVP

FOR AGRICULTURAL/COMMERCIAL USE ONLY. NOT FOR SALE FOR ANY OTHER USES.

KEEP JUT OF REACH OF CHILDREN CAUTION

See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENTS:

Dichlorvos: 2,2-Dichlorcvinyl dimethyl phosphate	.23%
Related compounds	.02%
Piperonyl butoxide, technical*	. 26%
Pyrethrins	.03%
INERT INGREDIENTS**	99.46%
Total	100.00%
	•

*Equivalent to 0.21% (butylcarbityl)(6-propylpiperonyl) ether and 0.05% related compounds. **Contains Mineral Seal Oil.

ACCEPTED

Net Contents: 2 1/2 Gallons

MAR 3 0 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Règ. No. 1386-418 EPA Est. No. 1386-0H-1



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid inhalation of vapor or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate water, feed or food-stuffs.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Contact a physician. See "Physician's Note," below. If Inhaled: Remove victim to fresh air. Apply respiration if indicated. If On Skin: Remove contaminated clothing and wash contaminated skin with soap and water. If In Eyes: Flush eyes with plenty of water. Get medical attention if imitation persists.

Physician's Note: Atropine is antidotal.

For Medical Emergency information call 1-800-228-5635, extension 138.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from animals being treated. Do not contaminate water by disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT! Do not overapply. Do not make direct application livestock more frequently than once per day. of more than the recommended rate animal or use of coarse sprays may wet the hide and cause injury. a fine spray mist at a distance of 2-3 feet as directed to deposit a light mist Do not apply when animals on the hair. are lathered or temperatures exceed 80° F.

PY-VONA Stock Fly Spray contains DDVP, piperonyl butoxide and pyrethrins in a mineral seal oil base. It is a ready-to-use spray for application directly to dairy cattle, beef cattle, hogs, sheep, goats, and horses to control house flies, horn flies, stable flies, and mosquitoes and to aid in control of face flies.

DIRECT SPRAYING OF LIVESTOCK — Apply as a finely atomized spray uniformly distributed over each animal at a rate of 1 to 2 fluid ounces per animal on dairy cattle, beef cattle, and horses, and at a rate of 1 fluid ounce per animal on hogs, sheep and goats. Reduce application rate by one-half for young animals.

Use good spray equipment to spray each animal, preferably from the rear, directing the spray along both sides, including the belly, flanks, and withers, down each leg, over the back and around the poll. Hold nozzle of sprayer 2 or 3 feet away from the animal, so as to deposit droplets on the hair. DO NOT WET THE HIDE. For milking dairy animals, apply the spray 20 to 30 minutes before beginning to milk, or after milking has been completed. DO NOT SPRAY UDDERS OF LACTATING DAIRY ANIMALS. DO NOT APPLY IN EXCESS OF RATES STATED PER ANIMAL. Repeat daily as needed. Do not apply regularly to calves under six months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates. Do not treat animals within one day of slaughter. Do not apply dermalty in combination with Trichlorfon.

BARN SPRAYING — To control house flies in dairy barns, milk rooms, loafing sheds, livestock barns, stables, and poultry houses, apply 2 fluid ounces per 1,000 cubic feet as a finely atomized spray, uniformly distributed throughout the area. To reduce air movement as much as possible, it is preferable to close windows and doors before spraying; leave them closed for 10 minutes after spraying and then ventilate the area. When used in enclosed areas do not remain in treated areas after spraying. Ventilate the area before reentering. Repeat daily, morning and afternoon, as needed. Do not apply as a barn spray if animals which have received direct application within 8 hours remain in the barn.

DO NOT CONTAMINATE FEED, DRINKING WATER, MILK, AND MILKING EQUIPMENT.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

CO-OP is a Trademark of Universal Cooperatives, Inc.

Manufactured by: Universal Cooperatives, Inc. Minneapolis, MN 55440

