



# 5.6% MANEB DUST

## INGREDIENTS

ACTIVE INGREDIENT:	
Maneb (Manganese ethylenebisdithiocarbamate) .....	5.6%
INERT INGREDIENTS .....	94.4%
Total .....	100.0%

EPA Reg. No. 1386-427

**CAUTION**  
Keep Out Of Reach Of Children

May be absorbed through skin and throat. May cause irritation of eyes, nose, throat, and skin. Do not breathe dust. In case of contact with eyes or skin, flush with plenty of water; for eyes, get medical attention. Avoid contamination of feed and foodstuffs. Keep container closed when not in use.

This product is poisonous to fish. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit located well away from water supplies.

**DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.**

## DIRECTIONS FOR USE

**TOMATOES**—To control early blight, late blight, Septoria leaf spot, gray leaf spot, and anthracnose on field grown tomatoes, apply uniformly and thoroughly to foliage and fruit at the rate of 30 to 40 pounds per acre. Begin applications before diseases appear, or when disease is first reported in the area, and repeat the 7 to 10-day intervals or oftener under severe disease conditions.

**POTATOES**—To control early blight and late blight, apply thoroughly to the foliage at the rate of 40 to 50 pounds per acre. Begin applications when plants are small (2 to 4 inches high), and repeat at 5 to 10-day intervals as long as blight conditions persist. Under severe disease conditions, use the shorter interval between applications.

**CUCUMBERS, MELONS, AND SQUASH**—To control downy mildew and anthracnose, apply thoroughly and uniformly at the rate of 30 to 40 pounds per acre. Begin applications when the runners start to form, or when disease is first reported in the area, and repeat at 7 to 10-day intervals. Under severe disease conditions, apply at 3 to 5-day intervals.

## NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

**NET WEIGHT**

**50 POUNDS**

FC-784-A-572

PROD. 1231

DISTRIBUTED BY  
**UNITED CO-OPERATIVES, INC**  
ALLIANCE OHIO 4316

AC  
July 10-1972  
UNDER  
FUNGICIDE  
OR EST.  
DUST  
1386-427