

5.6% MANEB DUST

KEEP OUT OF REACH OF CHILDREN

CAUTION


See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENT:


MANEB (Manganese ethylene bisdithiocarbamate).....	5.6%
INERT INGREDIENTS	94.4%
Total	100.0%

Net Weight: 50 Pounds
Prod. 1231

EPA Reg. No. 1386-427
EPA Est. No. _____



UNIVERSAL COOPERATIVES, INC.
MINNEAPOLIS, MN 55420



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Do not contaminate feed and foodstuffs. Protective clothing (long pants, long sleeve shirt, gloves, hat, and boots) must be worn during loading.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or POISON CONTROL CENTER immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. If inhaled: Remove victim to fresh air. Apply respiration if indicated. If on skin: Wash with soap and water. If in eyes: Flush eyes for at least 15 minutes with water. Call a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift of spray from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TOMATOES - To control early blight, late blight, Septoria leaf spot, gray leaf spot, and anthracnose on field grown tomatoes, apply uniformly and thoroughly to foliage and fruit at the rate of 30 to 40 pounds per acre. Begin applications before diseases appear, or when disease is first reported in the area, and repeat at 7 to 10-day intervals or oftener under severe disease conditions.

POTATOES - To control early blight and late blight, apply thoroughly to the foliage at the rate of 40 to 50 pounds per acre. Begin applications when plants are small (2 to 4 inches high), and repeat at 5 to 10-day intervals as long as blight conditions persist. Under severe disease conditions, use the shorter interval between applications.

CUCUMBERS, MELONS, AND SQUASH - To control downy mildew and anthracnose, apply thoroughly and uniformly at the rate of 30 to 40 pounds per acre. Begin applications when the runners start to form, or when disease is first reported in the area, and repeat at 7 to 10-day intervals. Under severe disease conditions, apply at 3 to 5-day intervals.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to approved Federal, state or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application

equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions, or warnings stated on this label.