




20% CHLORDANE GRANULAR



INGREDIENTS

ACTIVE INGREDIENTS:	
Technical chlordane*	20%
INERT INGREDIENTS	80%
Total	100%

*Equivalent to 12% octachloro-4,7-methanotetrahydroindane and 8% related compounds.
EPA Reg. No. 1386-435

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust or vapor. In case of contact with skin, wash with soap and water. Do not contaminate feed and foodstuffs.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is poisonous. Stay well away from the smoke.

DIRECTIONS FOR USE

PREPLANTING SOIL APPLICATIONS — DIRECT APPLICATION TO SOIL BROADCAST TREATMENTS

Apply as a uniform broadcast treatment of the surface of the soil (previously prepared for planting) at the rate per acre indicated, promptly follow with cultivation to mix the granules into the soil to a depth of 6 to 8 inches, preferably 7 to 14 days prior to planting or transplanting of the crops shown below. Properly applied broadcast preplanting soil treatments may be effective for more than one year; repeat applications as new infestations develop, but not more than once yearly. Use the amount shown after each crop to control the insects named.

CORN—To control corn rootworms and seed corn maggot, 15 to 20 pounds per acre. To control wireworms and white grubs, 25 to 40 pounds per acre.

TOBACCO—To control wireworms, white grubs, and green June beetle larvae, 20 pounds per acre.

POTATOES—To control wireworms and white grubs, 25 to 40 pounds per acre.

STRAWBERRIES—To control wireworms, 25 to 40 pounds per acre; to control white grubs, 50 pounds per acre.

IN-ROW TREATMENTS

Apply uniformly in a band 10 to 12 inches wide centered over the row, promptly work the insecticide into the top 6 inches of soil, preferably 7 days prior to planting and transplanting of the crops named below. In-row treatments should be repeated annually. Use the amount shown after each crop to control the insect named.

CORN—To control corn rootworms, and seed corn maggot, 10 pounds per acre. To control wireworms and white grubs, 12½ to 20 pounds per acre.

TOBACCO—To control wireworms and white grubs, 12½ to 20 pounds per acre.

POTATOES—To control wireworms and white grubs, 12½ to 20 pounds per acre.

STRAWBERRIES—To control wireworms, 10 pounds per acre; to control white grubs, 10 pounds per acre. Do not apply when fruit is present.

REFERENCE TO WIREWORMS IN ABOVE SECTIONS DOES NOT INCLUDE CHLORDANE-RESISTANT WIREWORMS AND SOUTHERN POTATO WIREWORMS.

ORNAMENTAL TURF AND LAWNS

To control white grubs, Japanese beetle larvae and green June beetle larvae, apply 25 to 40 pounds per acre uniformly distributed over the grass area and immediately sprinkle or drench lawn to wash insecticide into the soil. Apply when insect damage is first noted, preferably in late summer or early fall when grubs and larvae are small.

Do not apply to lawns or turf areas when people or animals are on them, and do not allow drift of dust from treated area. Keep children and pets off treated areas until insecticide has been washed into the soil and the grass has completely dried. Do not graze or feed forage from treated areas to livestock.



CHLORDANE DUST BASE

INGREDIENTS

ACTIVE INGREDIENTS:
Technical chlordane*
INERT INGREDIENTS

40%
60%

TOTAL

100%

*Equivalent to 24% octachloro-4,7-methanotetrahydroindane
and 16% related compounds
USDA Reg. No. 1386-496

FOR MANUFACTURING USE ONLY

(Sold only for the preparation of insecticide dust mixtures)

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish and wildlife. Manufacturing operations must include procedures which will prevent contamination of any body of water, through disposal of containers, wastes, rinse solutions, and unused chemical. Finished dust mixtures containing this product must be adequately labeled with instructions to prevent contamination of any body of water or the environment of birds and other wildlife.

Finished dust mixtures containing this product may be used only for the purposes for which official tolerances for chlordane have been established, within the limitations which permit compliance with these tolerances, and must be labeled with acceptable directions for use.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Completely burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

NET WEIGHT

50 POUNDS

FC 250-A467

PROG 1024

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO