



LICE-X[®]

GRANULAR INSECTICIDE

CONTROLS HOG LICE
On All Ages of Swine—Newborn, Too
Contains KORLAN[®]

INGREDIENTS

ACTIVE INGREDIENTS:	Ronnel [0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate]*	3%
INERT INGREDIENTS	95%
Total	100%

*From Technical Grade—Not Suitable for Internal Use in Livestock.
EPA Reg. No. 1386-448

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust or vapor. Avoid contact with skin. Wash thoroughly after using. Do not get in eyes. Avoid contamination of feed and foodstuffs.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

Apply the granules evenly over the bedding for swine at the rate of ½ pound per 100 square feet of area. The granules may be applied with a scoop or a can. Mixing with the bedding is not necessary. One treatment will control the hog-lice infestation when the animals use the treated bedding regularly.

Replace the treated bedding with clean, untreated bedding or otherwise exclude the swine from the treated bedding two weeks prior to slaughter.

LIVESTOCK REMEDY — NOT FOR HUMAN USE.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

! KORLAN[®] Proprietary name for ronnel. Registered trademark of The Dow Chemical Co.
U.S. Pat. No. 2,599,516

NET WEIGHT

5 POUNDS

FC-815-D472

PROD. 933

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

