

faces, such as walls, fences, garbage cans, and refuse containers, with 1 pint in 3 gallons of water (2-2/3 tbsp. in 1 quart). Repeat as necessary.

DO NOT CONTAMINATE FEED AND FOODSTUFFS, DRINKING FOUNTAINS, FEED TROUGHS, AND LITTER. DO NOT USE IN MILK-PROCESSING ROOMS, MILK HOUSES, AND MILK STORAGE ROOMS. DO NOT USE IN HOMES.

#### CAUTION

VAPOR HARMFUL. HARMFUL IF SWALLOWED. Concentrated material causes eye irritation. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with skin or eyes. Wash thoroughly after handling. Do not contaminate feed and foodstuffs. Keep away from heat, sparks and open flame.

In case of contact with eyes, flush with plenty of water for at least 15 minutes; get medical attention.

NOTE TO PHYSICIANS—This product, upon repeated, prolonged use, causes cholinesterase inhibitor. Atropine is antidotal.

Do not store where temperature may be below 32°F. Do not use this product for any purpose other than as directed in this label and accompanying leaflet.

Destroy empty containers. Do not reuse this container for any purpose. Break empty glass containers and bury fragments in an isolated area.

#### NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

USDA Reg. No. 1386-449

## CYGON 2-E SYSTEMIC INSECTICIDE

### CONTROLS HOUSEFLIES

#### ACTIVE INGREDIENT:

Dimethoate [0,0-dimethyl S-(N-methyl-carbamoylmethyl) phosphorodithioate] .... 23.4%

INERT INGREDIENTS ..... 76.6%

Total ..... 100.0%

CONTAINS 2 POUNDS DIMETHOATE PER GALLON

Directions for use on Ornamental Plants in  
Accompanying Leaflet

### CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

\*Trademark of American Cyanamid Co., Wayne, N.J.,  
U.S. Pat. 2,494,283

NET VOLUME 1 U.S. PINT LIQUID

FC-826E-1-67 DISTRIBUTED BY PROD. 259

## CYGON\* 2-E SYSTEMIC INSECTICIDE

### DIRECTIONS FOR USE

#### Control of Houseflies

**RESIDUAL SPRAY**—To control houseflies in dairy barns, hog pens, calf barns, loafing sheds, poultry houses, and other farm outbuildings, apply a 1 percent residual spray to ceilings, walls, stanchions, and other surfaces on which flies congregate or rest. Prepare the spray by mixing 1 gallon in 25 gallons of water (1 pt. in 3 gal.). Apply the spray to wet uniformly all fly-resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 sq. ft. of wall or ceiling surface. Repeat applications as necessary.

Remove dairy animals, calves under one month of age, and poultry from buildings when applying residual sprays.

**SPOT SPRAY**—To control houseflies in localized areas, apply a spray made by mixing 1/2 pint in 5 quarts of water (3 1/4 tbsps. in 1 qt.) to surfaces around doorways and windows, in alleyways and feed rooms, and to other spots frequented by flies. Repeat as necessary.

GOOD SANITATION IS A NECESSARY PART OF ANY EFFECTIVE FLY CONTROL PROGRAM.

**MAGGOT SPRAY**—To control housefly maggots, mix 1/2 pint in 5 quarts of water (1 qt. in 5 gal.) and apply as a coarse spray or with a sprinkling can to fly-breeding media, such as poultry droppings in caged layer houses, garbage dumps, or manure piles. Repeat applications as additional manure or garbage is added.

**OUTSIDE SPRAY**—To control houseflies around home, and recreation areas, animal quarters, food processing plants, warehouses, loading docks, and refuse areas, thoroughly spray exposed sur-

UNITED CO OPERATIVES INC

4/27/67

1386-449