



5% SEVIN® DUST

INGREDIENTS

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate)

INERT INGREDIENTS


TOTAL

5%
95%
100%

Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate

EPA Reg. No. 1386-451
CAUTION

See Remainder of CAUTION Statement on Back Panel



NET WEIGHT

FC-831-B-672

5 POUNDS

Prod. 1502

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

5% SEVIN® DUST

DIRECTIONS FOR USE

VEGETABLES AND SMALL FRUITS

For control of insects on the crops listed below apply at the rate stated in pounds per acre uniformly and evenly. After 1 application, or after 3 days of the single dust coverage of foliage should be complete. If 2 or more applications are made, the 2nd application should be applied 7 days later. Repeat as necessary. Generally 2 to 3 day intervals unless shorter interval is stated. May be applied on day of harvest if otherwise directed.

ASPARAGUS—African beetle, 40 lb. per acre. Do not repeat application within 3 days. Do not apply to the foliage until after cutting season.

BEANS—Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetles, Japanese beetle, adult and larva, 40 lb. per acre.

CORN—Sawfly, Corn earworm, European corn borer, fall armyworm, flea beetles, Japanese beetle, adult and larva, and leafhoppers, 40 lb. per acre. For larvae in whorls (European corn borer and fall armyworm) and earworms, apply to entire plant before tasseling. For insects attacking silk and ear, apply at 2 to 3 day intervals starting when silk first appear and continuing until they become dry. Three or more applications may be required. Proper timing and thorough coverage are essential. Application of this product to the tassel stage of corn during pollen shed period will seriously reduce bee population.

CUCUMBERS, MELONS, PUMPKINS, AND SQUASH—Pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers and squash bug, 20 lbs. per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when foliage is wet or during period of prolonged high humidity. Do not use on watermelon.

POTATOES, TOMATOES, EGGPLANT, AND PEPPERS—Colorado potato beetle, flea beetles, leafhopper, Colorado potato beetle, tomato worm, tomato fruitworm, tomato hornworm, tarnished plant bug, and stink bug, 40 lb. per acre.

CABBAGE, BROCCOLI, AND CAULIFLOWER—Flea beetles, Harlequin bug, imported cabbage worm, and imported cabbageworm, 40 lb. per acre. Do not apply within 3 days of harvest.

GARDEN BEETS, LETTUCE, RADISHES, SPINACH, TURNIPS AND CARROTS—Flea beetles, Harlequin bug, leafhoppers, corn rootworm, cornworm, picklebug, tarnished plant bug, and stink bug, 40 lb. per acre. Do not apply to garden beets, head lettuce, radishes, and turnips within 3 days of harvest. Do not apply to leaf lettuce and spinach or to garden beets and turnips if tops are to be used for food or feed within 14 days of harvest.

STRAWBERRIES—Meadow spittlebug, strawberry leaf roller, and strawberry weevil, 40 lb. per acre. Do not apply within 1 day of harvest.

GRAPE—Grape berry moth, Japanese beetle adults, red-handed leaf roller, leafhoppers, and grape leaf weevils, 40 lb. per acre.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, AND RASPBERRIES—Japanese beetle adults, leafhoppers, and aphid, 40 lb. per acre. Do not apply within 7 days of harvest.

BLUEBERRIES—Cherry maggot, cherry fruitworm, and cranberry fruitworm, 40 lb. per acre. Applications to blueberry mound should be timed with emergence of adult flies and repeated according to local schedules.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions, or warnings stated on this label.

ORNAMENTAL PLANTS

FLOWERS, SHRUBS AND SHADE TREES—Apple aphid, bagworm, birch leaf miner, box elder bug, boxwood leaf miner, elm leaf beetle, Japanese beetle adults, lace bugs, leafhoppers, leaf roller, mealy bug, mimosa webworm, pine needle scale, rose aphid, rose slug, tent caterpillar, and thrips (excepted on ornamental plants and shade trees such as rose, azalea, chrysanthemum, gladiolus, lilac, arbor vitae, juniper, hydrangea, azalea, elm, rose, oak, dogwood, birch and pine). Apply thoroughly and uniformly to upper and lower surfaces of all foliage. Reapply at 7-day intervals as needed. Do not apply to Boston ivy.

POULTRY

CHICKENS, PARTRIDGES, PHEASANTS, AND PIGEONS—Northern fowl mite, chicken red mite, lice, fleas, and feather mites, 1 pound. Apply 1 pound per 40 q ft of floor and litter, and to nest and exterior feathers. Do not apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of application.

For 100 broiler chickens, add 1 lb. of the birds, as a supplement to treatment of premises, apply 1 pound per 100 q ft. Divide among 100 chickens, and distribute throughout feathers. Do not repeat within 4 weeks. Do not use on 7 day old chicks.

DO NOT CONTAMINATE FEED, DRINKING WATER, FEEDING AND WATERING RECEPTACLES

PET ANIMALS

DOGS AND CATS—Tick and flea. Apply liberally over entire animal avoiding eyes, and rub thoroughly to insure that flea life cycle is destroyed after treatment. Repeat at weekly intervals as needed. Do not apply to cats less than 4 weeks old.

Flea and tick control may be aided by feeding cats. Apply liberally into oil around base of neck, between shoulder blades, and creases of loom, and belly. Repeat as necessary.

DO NOT CONTAMINATE FEED OR FEEDING AND WATERING RECEPTACLES

SPECIAL NOTES

Do not apply to foliage on tender foliage in the presence of rain or high humidity of 70% or more. Do not apply to foliage containing the soft scale spider mites, but is compatible with hot commodity dust materials. Do not apply to foliage of tobacco under 4 weeks of age. This product is highly toxic to the earthworm. Direct contact with the skin may cause irritation. Protective equipment may be obtained from your distributor. Agreed to by Executive Order.

CAUTION

DO NOT EAT, DRINK, OR SMOKE. Do not breathe dust. Do not take internally. Skin contact may be painful. Avoid contact with eyes. Wash hands before eating. Take shower or bath after work. Wear rubber long-sleeved shirt and pants. Change clothes daily. Do not contaminate feed and foodstuffs.

OTC FOP-PHYS-A—Carcinogen, a moderate reversible cholesterolase inhibitor. At present intended for use in humans only. Keep out of any body of water. Apply only for the purpose and in the manner indicated on label. Do not spray when weather conditions favor drift of dust from treated surface. Insecticide and fungicide, or other chemical in a pit on non-crop land well away from water supplies.

DO NOT STORE OUTDOORS—The make may be hazardous. Store in a cool, dry place.

