

**ALFALFA
WEEVIL SPRAY**

Controls Alfalfa Weevil Larvae, Aphids, Leafhoppers,
Spittlebug, and Clover Leaf Weevil on Alfalfa and Clover.

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel For Additional Precautionary Statements


ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	23.2%
Methoxychlor, technical*	23.2%
Xylene	46.7%
INERT INGREDIENTS	6.9%
Total	100.0%

*Equivalent to 20.4% 2,2-Bis(p-methoxyphenol)-1,1,
1-trichloroethane and 2.8% other isomers and related
compounds. Contains 2 pounds malathion and 2 pounds
technical methoxychlor per gallon.


NET CONTENTS: 1 GALLON
PROD. 262

EPA Reg. No. 1386-464
EPA Est. No. 1386-OH-1



UNIVERSAL COOPERATIVES, INC.
MINNEAPOLIS, MN 55440

ALSO AVAILABLE IN...



LE

PRELIMINARY STATEMENTS
REGARDS TO HUMAN AND DOMESTIC ANIMALS
WARNING

Causes eye irritation. Do not get in eyes, on skin or on clothing. Be careful if swallowed. Wash thoroughly after using. Avoid contact with food and feedstuffs.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician or Poison Control Center. Do not induce vomiting because of possible aspiration hazard. Get medical attention. If on skin, wash affected areas with soap and water. If inhaled, remove victim to fresh air. Apply respiration if indicated. If in eyes, flush eyes for at least 15 minutes with water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALFALEA - To control alfalfa weevil larvae, use 2 to 3 quarts (1 to 1.5 lb. malathion and 1 to 1.5 lb. methoxychlor) per acre in the amount of water required for uniformly thorough spray coverage. Begin applications when larvae appear, usually in May or June, and before they cause serious damage. Repeat at 10 to 14-day intervals as necessary. In some areas, it may be necessary to spray the stubble after removal of the first cutting.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR FEEDING.

DO NOT APPLY DURING FULL BLOOM.

CLOVER AND ALFALEA - To control aphids, spittlebugs, leafhoppers, clover leaf weevil, alfalfa caterpillar, and grasshoppers, use 2 to 3 quarts per acre in the amount of water required for uniformly thorough spray coverage. Begin applications when insects first appear and repeat at 7 to 10-day intervals as necessary.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR FEEDING. DO NOT APPLY DURING FULL BLOOM.

The amount of water required for uniformly thorough spray coverage is usually 15 to 20 gallons per acre for boom sprayers, although as little as 10 gallons per acre has been used successfully in some instances.

MR. L. C. WILLIAMS
Universal Cooperative, Inc.
3001 Metro Drive, Suite 500
Minneapolis, MN 55420

Dear Mr. Williams:

Subject: Alfalfa Weevil Spray
EPA Registration No. 1380-464
Your Application Dated August 19, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (75-767)

Enclosure

TS-767;ELLENBERGER;DCR-24748;WANG-0319C;CBP;Raven:479-2013;9/23/82