

such as heat, except for spot spraying. Most legumes, including clover, are severely damaged, and under some conditions, killed.

Do not contaminate domestic or irrigation water supplies.

Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination and growth.

Do not graze or feed forage from treated ones to dairy animals within 7 days after treatment.

It is almost impossible to remove residues of 2,4-D from sprayers and spray equipment, particularly from non-metallic parts, e.g., wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

ACCEPTED
5/10/67
1386 102

Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, trees, fruit trees, ornamentals, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of this product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigating water supplies.

It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g. wood, rubber, polyethylene fibre, and the equipment used to measure and apply this weed killer should be used subsequently for applying other materials to plants or crops. Destroy containers; never re-use for any purpose.

This product is poisonous to fish. Keep out of any body of water for the purposes and in the manner stated in this label. Do not apply under conditions favor drift of spray from area treated. Clean application equipment, dispose of rinse solutions, wastes, and unused chemical in a safe manner, well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is plastic, punch holes in top and bottom, crush under wheels of heavy vehicle or with an iron pipe, and bury crushed container in a pit on non-crop land well away from water supplies. Do not burn empty containers.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.
BANVEL (R) D-T.M. of Velsicol Chem. Corp.



INGREDIENTS:

2,4-Dichlorophenoxyacetic acid, diethylethanolamine salt*	19.9%
Licamba (2-methoxy-3,6-dichlorobenzoic acid, dimethylamine salt)**	6.6%
INERT INGREDIENTS	73.5%
Total	100.0%

*Equivalent to 16.5% 2,4-dichlorophenoxyacetic acid
**Equivalent to 5.53% 2-methoxy-3,6-dichlorobenzoic acid
CONTAINS 1.5 LB. 2,4-D ACID EQUIVALENT AND 0.5 LB. DICAMBA PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 1386-467

Net Volume 1 U.S. Quart Liquid

H&G-122C-2-72 MANUFACTURED BY PROD. 2118

UNICO Turf Treeter Lawn Weed Killer

Contains BANVEL (R) D and 2,4-D

For Chickweed, Knotweed, Red Sorrel, and Many Common Broadleaf Weeds in Lawns and Ornamental Turf

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF—To control chickweeds (common, mouse-ear and perennial or stitchwort), sheep sorrel, knotweed, clover, knawel, curly dock, dog fennel, chickory, mustard, ragweed, lambsquarter, plantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, and stinging nettle in established perennial grasses in lawns or ornamental turf, use 3 fluid ounces (6 tablespoonfuls) in 1 gallon of water and apply this mixture as a spray to wet uniformly the weed plants on 1,000 square feet (20' x 50'). This rate of application is equivalent to 1 gallon of product per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition.

Do not apply on or near desirable trees, shrubs, and other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not clean application equipment or utensils in these areas. Do not apply to centipede grass, St. Augustine grass, or to dichondra. Do not use on lawns where desirable bent grasses or clovers are present.

DO NOT GRAZE TREATED AREAS OR FEED TREATED FORAGE OR THRESHINGS TO LIVESTOCK. DO NOT APPLY TO NEWLY SEEDED LAWNS OR TURF.

CAUTION

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Do not store with or near fertilizers, seeds, insecticides, or fungicides. Do not use or store near heat or open flame.

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamentals, corn or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of this product may cause volatilization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigation water supplies.

It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g. wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops. Destroy empty containers: never re-use for any purpose.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies. Do not burn empty containers.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.
BANVEL (R) D-1-M of Velicol Chem. Corp



ACTIVE INGREDIENTS:	
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	19.9%
Dicamba (3-methoxy-2,4-dichlorobenzoic acid, dimethylamine salt)**	6.6%
INERT INGREDIENTS	73.5%
Total	100.0%

*Equivalent to 16.5% 2,4-Dichlorophenoxyacetic acid
**Equivalent to 5.50% 3-methoxy-2,4-dichlorobenzoic acid
CONTAINS 1.5 LB. 2,4-D ACID EQUIVALENT AND 0.5 LB. DICAMBA PER GALLON

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 188A-07

Net Volume 1 U.S. Quart Liquid
MFG. 1-28-72 MANUFACTURED BY PRGO. 2118

Do not apply to areas where there may be exposed water supplies. Do not use on fields where there may be exposed water supplies. Do not use on fields where there may be exposed water supplies. Do not use on fields where there may be exposed water supplies.

UNITED CO OPERATIVES, INC.
ALLIANCE OHIO 44601

6-13-72
1386-467

21

increased in using this product to avoid injury to susceptible flowers, grapes, fruit trees, ornamental plants and do not permit spray mist to drift onto susceptible plants during both cool and hot weather. Do not apply during both cool and hot weather. Do not apply during both cool and hot weather. Do not apply during both cool and hot weather.

residues of this product from sprayers and spray nozzles, e.g. wood, rubber, polyethylene, or other plastic. Do not apply this weed killer should not be used on plants of crops. Destroy empty containers.

Keep out of any body of water. Apply only as directed in this label. Do not apply when weather is hot or humid. Clean application equipment and store in a safe place in a pit on non-crop land.

Do not reuse this container for any purpose. If reused, thoroughly clean and dry. If container is metal, punch holes in it with a sharp object or with an axe, and bury it well away from water supplies. Do not reuse.

NOTICE
Read and comply with the limitations, cautions or directions on this label. For more information, contact the manufacturer.



ACTIVE INGREDIENTS:	
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	15.9%
Dicamba (2-methoxy-3,6-dichlorobenzoic acid, dimethylamine salt)**	6.6%
INERT INGREDIENTS	73.5%
Total	100.0%
*Equivalent to 16.5% 2,4-dichlorophenoxyacetic acid	
**Equivalent to 5.53% 2-methoxy-3,6-dichlorobenzoic acid	
CONTAINS 1.5 LB. 2,4-D ACID EQUIVALENT AND 0.5 LB. DICAMBA PER GALLON	

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 1306-057
Net Volume 1 U.S. Quart Liquid
NAB-128C-2-72 MANUFACTURED BY PROD 2118

UNICO Turf Treeter Lawn Weed Killer

Contains 2,4-D and Dicamba

For Chickweed, Knotweed, Red Sorrel, and Many Common Broadleaf Weeds in Lawns and Ornamental Turf

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF: To control chickweeds (common, mouse-ear and perennial or stitchwort), knotweed, red sorrel, lambsquarters, plantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, and stinging nettle in established perennial grasses in lawns or ornamental turf, use 3 fluid ounces (6 tablespoons) in 1 gallon of water and apply this mixture as a spray to wet uniformly the weed plants on 1,000 square feet (20' x 50'). This rate of application is equivalent to 1 gallon of product per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition.

Do not apply on or near desirable trees, shrubs, and other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not clean application equipment or storage in these areas. Do not apply to centipede grass, St. Augustine grass, or to dicamba. Do not use on lawns where desirable warm grasses or clovers are present.

DO NOT GRAZE TREATED AREAS OR FEED TREATED FORAGE OR THRESHING TO LIVESTOCK. DO NOT APPLY TO NEWLY SEEDED LAWNS OR TURF.

CAUTION

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Do not store with or near fertilizers, seeds, insecticides, or fungicides. Do not use or store near heat or open flame.

214

UNITED CO OPERATIVES, INC
ALLIANCE OHIO 44601

RECEIVED
6-13-72
1386-467

21



Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of this product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigating water supplies.

It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g. wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops. Destroy empty containers; never re-use for any purpose.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies. Do not burn empty containers.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.
BANVEL (R) ©-T.M. of Velsicol Chem. Corp.



**TURF TREETER
LAWN WEED KILLER**

ACTIVE INGREDIENTS:	
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	19.9%
Dicamba (2-methoxy-3,6-dichlorobenzoic acid, dimethylamine salt)**	6.6%
INERT INGREDIENTS	73.5%
Total	100.0%

*Equivalent to 16.5% 2,4-dichlorophenoxyacetic acid
**Equivalent to 5.53% 2-methoxy-3,6-dichlorobenzoic acid
CONTAINS 1.5 LB. 2,4-D ACID EQUIVALENT AND 0.5 LB. DICAMBA PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 1386-467

Net Volume 1 U.S. Quart Liquid

H&G-122C-2-72 MANUFACTURED BY PROD. 2118

**UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601**

UNICO Turf Treeter La

Contains BANVEL (R) D

For Chickweed, Knotweed, Red Sorrel, and Weeds in Lawns and Orna

DIRECTIONS FOR

LAWNS AND ORNAMENTAL TURF—To control and perennial or stitchwort), sheep sorrel, knotweed, fennel, chickory, mustard, ragweed, lambsquarter, wild radish, bull thistle, burdock, dandelion, and perennial grasses in lawns or ornamental turf, use 3 1 gallon of water and apply this mixture as a spray on 1,000 square feet (20' x 50'). This rate of application per acre. Apply in the spring or early summer, rapidly growing condition.

Do not apply on or near desirable trees, shrub where their roots may extend, or in locations where moved into contact with their roots. Do not clean in these areas. Do not apply to centipede grass, St. Do not use on lawns where desirable bent grasses or

DO NOT GRAZE TREATED AREAS OR FEED TRE TO LIVESTOCK. DO NOT APPLY TO NEWLY SEED

CAUTION

Harmful if swallowed. Do not take internally, and clothing. In case of contact, flush with plenty of water. Do not store with or near fertilizers, seeds not use or store near heat or open flame.

6-13-72
1386-467

21

21A



UNICO Turf Treeter Lawn Weed Killer

Contains BANVEL (R) D and 2,4-D

For Chickweed, Knotweed, Red Sorrel, and Many Common Broadleaf Weeds in Lawns and Ornamental Turf

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF—To control chickweeds (common, mouse-ear and perennial or stitchwort), sheep sorrel, knotweed, clover, knawel, curly dock, dog fennel, chickory, mustard, ragweed, lambsquarters, plantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, and stinging nettle in established perennial grasses in lawns or ornamental turf, use 3 fluid ounces (6 tablespoonfuls) in 1 gallon of water and apply this mixture as a spray to wet uniformly the weed plants on 1,000 square feet (20' x 50'). This rate of application is equivalent to 1 gallon of product per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition.

Do not apply on or near desirable trees, shrubs, and other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not clean application equipment or utensils in these areas. Do not apply to centipede grass, St. Augustine grass, or to dichondra. Do not use on lawns where desirable bent grasses or clovers are present.

DO NOT GRAZE TREATED AREAS OR FEED TREATED FORAGE OR THRESHINGS TO LIVESTOCK. DO NOT APPLY TO NEWLY SEEDS LAWNS OR TURF.

CAUTION

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Do not store with or near fertilizers, seeds, insecticides, or fungicides. Do not use or store near heat or open flame.

TURF TREETER LAWN WEED KILLER

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	19.9%
Dicamba (2-methoxy-3,6-dichlorobenzole acid, dimethylamine salt)**	6.6%

INERT INGREDIENTS

Total 73.5%

*Equivalent to 16.5% 2,4-dichlorophenoxyacetic acid
 **Equivalent to 5.53% 2-methoxy-3,6-dichlorobenzole acid
CONTAINS 1.5 LB. 2,4-D ACID EQUIVALENT AND 0.5 LB. DICAMBA PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 1386-467

Net Volume 1 U.S. Quart Liquid

H8G-122C-2-73 MANUFACTURED BY PROD. 2118

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

72
467

21

21A