

10-26-89

Thorobred "H-P"® Spray

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Remainder Of CAUTION Statement On Side Panel

ACTIVE INGREDIENTS:


Technical piperonyl butoxide*	1.0%
Pyrethrins1%
INERT INGREDIENTS	98.9%
Total	100.0%

*Equivalent to 0.8% (butylcarbityl) (6-propyl-piperonyl) ether and 0.2% related compounds. Technical piperonyl butoxide and pyrethrins are constituents of PYRENONE® - Trademark of Fairfield American Corporation.

Net Volume: 1 Quart

EPA Reg. No. 1386-480
 EPA Est. No. 1386-OH-1

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Products - Subsection for Phase 2
of FIFRA, Page 2.12, Article 15.14, and 15.15 in State.



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed, food-stuffs, water, watering devices, and feed troughs.

Do not wet the skin of animals as this may cause blistering. Do not apply to plants or vegetation. Do not store or use near neat or open flame. This product is intended for use on horses and ponies only. The directions are not adequate for any other purpose.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT! Do not overapply. Use of more than the recommended rate per animal or use of coarse sprays may wet the hide and cause injury. Use only a fine spray mist at a distance of 2-3 feet as directed to deposit a light mist on the hair. Do not apply when animals are lathered or temperatures exceed 80° F.

Thorobred "H-P" Spray should be applied in barn foggers and sprayers which dispense an atomized mist.

DIRECT SPRAYING OF HORSES—To protect horses against horn flies, houseflies, stable flies, and mosquitoes, apply 1 fluid ounce as a finely atomized spray uniformly distributed over each animal. Use a good hand sprayer or electric sprayer to spray each animal, preferably from the rear, directing the spray along the sides, including the belly, flanks, and withers, down each leg, over the back, and around the poll. Hold nozzle of sprayer 2 to 3 feet away from animal so as to deposit droplets on the hair. **DO NOT WET THE HIDE.** Repeat daily, morning, and afternoon, as needed.

BARN FOGGING—To control houseflies, stable flies, and horn flies, on the horses and in the barn, apply as a fog or finely

atomized mist over the horses in stable; use fogger or sprayer unit for each 2,500 sq. ft. of space above the animals and preferably near the ceiling. Doors and windows should be closed prior to application to prevent drafts and permit escape of the product until spray mist or fog is thoroughly distributed over the entire area; spray out 1 fl. oz. uniformly per animal. ventilate the barn for 15 minutes after the application. **REPEAT DAILY, MORNING AND AFTERNOON.** Keep sprayer, container, and nozzle clean. Do not deliver of fog or spray mist.

STORAGE AND DISPOSAL

STORAGE: Do not store near heat or open flame.

PESTICIDE DISPOSAL: Do not contaminate water by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL: Triple rinse and reuse. Empty containers should be disposed of in a sanitary landfill or by other means in accordance with local procedures.

WARRANTY AND LIMITATION

Seller warrants that this material conforms to the description and is reasonably fit for the purpose stated on the label when used in accordance with the directions of use and **Buyer** assumes the responsibility for any injury contrary to such directions. **Seller** makes no warranty, including any other express or implied warranty, of **Fitness** or of **Merchantability**, and is not authorized to do so except in writing with this warranty. In no event shall **Seller's** liability exceed the purchase price of the product if a claim is made.

