



# **Tomato-Potato & Vegetable DUST CONTAINS SEVIN®-COPPER**

## INGREDIENTS

## ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) . . . . .	5%
Copper, expressed as metallic (in copper oxide) . . . . .	7%
<b>INERT INGREDIENTS:</b> . . . . .	<b>88%</b>
Total	100%

EPA REG. NO. 1386-499

## **CAUTION**

KEEP OUT OF REACH OF CHILDREN

Harmful if INHALED or SWALLOWED; may cause irritation of skin, eyes, nose, throat or lungs; avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using; change clothing.

This product is poisonous to fish or wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated on this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies. Avoid contamination of food and feedstuffs.

**DESTROY EMPTY CONTAINERS** — Do not reuse this container for any purpose. Destroy container in compliance with local regulations if burned stay away from smoke smoke may be hazardous.

**BEEES** — Honeybees should be located a safe distance from all areas to be treated during the growing season. Avoid applications during periods when beneficial bees are visiting the plantings.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is not advised.

To control insects, plant diseases and field grown crops listed below, apply at the rate specified for each crop to upper and lower surfaces of the foliage. Dust coverage of 5 lbs/acre of 1 lb. bags and thorough, excessive dusting may scald tender foliage. Begin applications when the insects or their damage first appear and before diseases appear. When disease is first reported in the area, repeat as necessary, generally at 7 day intervals unless a shorter interval is stated. Some damage may occur on tender foliage in the presence of rain or high humidity of several days duration.

**TOMATOES** — Tomato fruitworm, tomato hornworm, leaf rollers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, ear, blight, late blight and anthracnose on field grown tomatoes. Apply 35 to 40 pounds of dust per acre. Do not apply same day as harvest.

ings; avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using; change clothing.

This product is poisonous to fish & wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated on this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes of unused chemical in a pit on non-crop land well away from water supplies. Avoid contamination of feed and foodstuffs.

**DESTROY EMPTY CONTAINERS** — Do not reuse the container for any purpose. Destroy immediately. Do not burn. In the event of a fire, stay away from smoke, smoke may be dangerous.

**BEES** — Honeybees should be located a safe distance from all areas to be treated during the growing season. Avoid applications during periods when beneficial bees are visiting the plant or crop.

**NOTE FOR PHYSICIAN** — Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

To control insect-caused diseases on field-grown crops listed below, apply at the rate specified for each crop to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough; excessive dusting may scald tender foliage. Begin applications when the insects or their damage first appear and before diseases appear when disease is first reported in the area. Repeat as necessary generally at 7-day intervals unless a shorter interval is stated. Some damage may occur on tender foliage in the presence of rain or high humidity if several days duration.

**TOMATOES** — Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall armyworm, corn borer, Colorado potato beetle, early blight, late blight, and anthracnose on field-grown tomatoes. Apply 35 to 40 pounds of dust per acre. Do not apply same day of harvest, etc.

**POTATOES** — Colorado potato beetle, leafhoppers, flea beetles, fall armyworm, European corn borer, early blight, late blight. Apply 35 to 40 pounds of dust per acre to foliage only. Do not apply same day of harvest.

**BEETS** — Flea beetles, spittlebug, tarnished plant bug, stink bug, cabbage looper (when larvae are small), aphid, 4 to 5-day meadowlark, and leaf spot. Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED. OTHERWISE NOT LATER THAN 2 DAYS OF HARVEST.**

**EGGPLANT AND PEPPERS** — Leafhoppers, fall armyworms, European corn borer, tarnished plant bug, and aphids. Apply 35 to 40 pounds of dust per acre. **DO NOT APPLY SAME DAY OF HARVEST.**

**CUCUMBERS, MELONS AND SQUASH** — Cucumber beetle, leafhoppers, squash bugs, aphid, fall armyworm, downy mildew, and anthracnose. Apply 20 pounds of dust per acre. Do not use on watermelons. **DO NOT APPLY SAME DAY OF HARVEST.**

**BEANS** — Mexican bean beetle, bean leaf beetle, flea beetle, leafhoppers, tarnished plant bug, and velvet bean caterpillar. Apply 40 pounds of dust per acre. **DO NOT APPLY SAME DAY OF HARVEST.**

**CORN** — Flea beetles, leafhoppers, corn rootworm adults (cucumber beetles), sap beetles, Japanese beetles, corn earworm, and European corn borer. Apply at rate of 40 pounds per acre. **DO NOT APPLY SAME DAY OF HARVEST; DO NOT FEED FORAGE TO DAIRY ANIMALS OR ANIMALS BEING FURNISHED FOR SLAUGHTER.**

**APPLICATION OF THIS PRODUCT TO THE TASSEL REGION OF CORN DURING THE POLLEN SHED PERIOD WILL SERIOUSLY REDUCE BEE POPULATION**

#### NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

**SEVIN®** — Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

**NET WEIGHT**

FC-898 A1271

**50 POUNDS**

PROD. 1540

**DISTRIBUTED BY**  
**UNITED CO-OPERATIVES, INC.**  
**ALLIANCE, OHIO 44601**