



Tomato-Potato & Vegetable DUST

CONTAINS SEVIN®-COPPER

INGREDIENTS:

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	5%
Copper, expressed as metallic (in copper oxide)	7%

INERT INGREDIENTS:	88%
--------------------	-----

Total	100%
-------	------

EPA REG. NO. 1386-499

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if INHALED or SWALLOWED; may cause irritation of skin, eyes, nose, throat or lungs; avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using; change clothing.

This product is poisonous to fish or wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated on this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies. Avoid contamination of feed and foodstuffs.

DESTROY EMPTY CONTAINERS. — Do not reuse this container for any purpose. Destroy container in compliance with local regulations; if burned stay away from smoke; smoke may be hazardous.

BEES — Honeybees should be located a safe distance from all areas to be treated during the growing season. Avoid applications during periods when beneficial bees are visiting the plants or crops.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

To control insects, plant diseases on field-grown crops listed below, apply at the rate specified for each crop, to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough, excessive dusting may scald tender foliage. Begin applications when the insects or their damage first appear and before diseases appear (when disease is first reported in the area). Repeat as necessary, generally at 7-day intervals unless a shorter interval is stated. Some damage may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES — Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, early blight, late blight, and anthracnose on field grown tomatoes. Apply 35 to 40 pounds of dust per acre. Do not apply same day of harvest.

POTATOES — Colorado potato beetle, leafhoppers, flea beetles, fall armyworm, European corn borer, early blight, and late blight. Apply 40 pounds of dust per acre to foliage only. Do not apply same day of harvest.

BEETS — Flea beetles, spittlebug, tarnished plant bug, stink bug, cabbage looper (when larvae

1-30-72
1386-499