


UNICO
Z Dust
Contains 5.2% Zineb Fungicide

INGREDIENTS

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bisdithiocarbamate) 5.2%
INERT INGREDIENTS. 94.8%
Total. 100.0%



1386-509

CAUTION (18-point type)
Keep Out of Reach of Children (12-point type)

See Remainder of CAUTION Statement on Back Panel


USDA Reg. No. 1386-359

Net Weight

5 Pounds

Distributed By
United Co-Operatives, Inc.
Alliance, Ohio

UNICO Z - Dust
Contains 5.2% Zineb Fungicide



CAUTION

Harmful if swallowed. May cause irritation of nose, throat, eyes, and skin. Do not breathe dust. In case of skin contact, wash thoroughly with soap and water. Avoid contamination of feed and foodstuffs. Store in a cool dry place. Keep container closed when not in use.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

Begin applications when plants are small or before symptoms of disease appear. Apply the dust at the rate of 30 to 40 pounds per acre, depending upon size of plants, as a uniformly thorough coverage of both upper and under surfaces of foliage. Repeat applications at 5 to 10-day intervals, depending upon the severity of diseases and on weather conditions which favor development of disease.

Use on the following crops or plants to control the diseases indicated.

TOMATOES - Early blight, late blight, anthracnose, Stemphylium and Septoria leaf spots, and Cladosporium gray mold. Begin applications when disease is first reported in area. Under severe blight conditions, repeat applications at 5 to 10-day intervals or oftener if necessary.

CELERY - Early blight and late blight - Begin dusting in the plant bed when plants first come through the ground. Make further applications at 3 to 5-day intervals until plants are set in the field. Continue field applications on regular weekly schedule. REMOVE RESIDUE BY STRIPPING, TRIMMING AND WASHING.

LETTUCE - Downy mildew, DO NOT APPLY TO HEAD LETTUCE WITHIN 5 DAYS OF

-3-

ACCEPTED
Nov 18 1970
1386-589

HARVEST OR TO LEAF LETTUCE WITHIN 7 DAYS OF HARVEST.

BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIPS - Cercospora leaf spot and downy mildew - DO NOT APPLY TO KALE, MUSTARD GREENS AND SWISS CHARD WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO BEETS AND TURNIPS WITHIN 7 DAYS OF HARVEST IF TOPS ARE TO BE USED FOR FOOD OR FEED.

CARROTS - Alternaria leaf blight and Cercospora leaf spot - Begin applications when diseases are reported in the area and continue at 7 to 10-day intervals during the growing season or as required. DO NOT APPLY TO CARROTS WITHIN 7 DAYS OF HARVEST IF TOPS ARE TO BE USED FOR FOOD OR FEED.

CUCUMBERS, MELONS, SQUASH - Anthracose, Alternaria (Macrosporium) blight, gummy stem blight, and downy mildew - Start applications when runners begin to form or at the first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves.

LIMA BEANS, DRY BEANS, STRING BEANS - Downy mildew, rust and Anthraconose - Apply when mildew first appears and repeat at weekly intervals. DO NOT APPLY TO BEANS WITHIN 7 DAYS OF HARVEST.

ONIONS - Downy mildew, blast (Botrytis leaf blight) and purple blotch - Begin applications when plants reach 4 leaf stage or when diseases are reported in the area and continue at 7 to 10-day intervals during the season. DO NOT APPLY TO GREEN ONIONS WITHIN 7 DAYS OF HARVEST.

SPINACH - Downy mildew (bluemold) and white rust - Apply at first sign of mildew and repeat at 7 to 10-day intervals or as required. DO NOT APPLY TO SPINACH WITHIN 7 DAYS OF HARVEST.

GRAPES - Downy mildew, black rot and ripe rot - Apply just before and just after bloom and again in 10 days. DO NOT APPLY TO GRAPES WITHIN 7 DAYS OF HARVEST.

STRAWBERRIES - Leaf spot - Begin when new growth starts in the spring. Repeat until fruit begins to form. Leaf scorch - Apply in new beds in late

Contd.

-4-

summer, making 2 to 3 weekly applications. DO NOT APPLY TO STRAWBERRIES WITHIN 7 DAYS OF HARVEST.

AZALEAS AND CAMELIAS - Petal blight - Apply 2 to 3 times a week while flowers are opening, and to the soil and litter around the plants while they are in blossom.

GIADIOLUS - Stemphylium leaf spot - Apply when disease appears and repeat every 7 to 10-days or oftener under severe disease conditions.

CARNATION - Leaf spot. DAHLIA - Botrytis. LILY - Botrytis. PANSY - Anthracose. ROSES - Black Spot. ZINNIA - leaf spot. Apply at weekly intervals through out the season.

NOTICE

Use only for the purpose and in compliance with the limitations cautions or warnings stated on this label.

ACCEPTED
Nov 18 1970
1386-589