

Harmful if **INHALED** or **SWALLOWED**; may cause irritation of skin, eyes, nose, throat or lungs; avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using; change clothing.

This product is poisonous to fish or wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated on this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies. Avoid contamination of feed and foodstuffs.

DESTROY EMPTY CONTAINERS. — Do not reuse this container for any purpose. Destroy container in compliance with local regulations; if burned stay away from smoke; smoke may be hazardous.

BEEES — Honeybees should be located a safe distance from all areas to be treated during the growing season. Avoid applications during periods when beneficial bees are visiting the plants or crops.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

To control insects, plant diseases on field-grown crops listed below, apply at the rate specified for each crop, to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough, excessive dusting may scald tender foliage. Begin applications when the insects or their damage first appear and before diseases appear (when disease is first reported in the area). Repeat as necessary, generally at 7-day intervals unless a shorter interval is stated. Some damage may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES — Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, early blight, late blight, and anthracnose on field grown tomatoes. Apply 35 to 40 pounds of dust per acre. Do not apply same day of harvest.

POTATOES — Colorado potato beetle, leafhoppers, flea beetles, fall armyworm, European corn borer, early blight, and late blight. Apply 40 pounds of dust per acre to foliage only. Do not apply same day of harvest.

BEETS — Flea beetles, spittiebug, tarnished plant bug, stink bug, cabbage looper (when larvae are small; apply at 4 to 5-day intervals), and leaf spot. Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED; OTHERWISE NOT WITHIN 3 DAYS OF HARVEST.**

EGGPLANT AND PEPPERS — Leafhoppers, fall armyworms, European corn borer, tarnished plant bug, and phomopsis. Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY SAME DAY OF HARVEST.**

CUCUMBERS, MELONS, AND SQUASH — Cucumber beetles, leafhoppers, squash bug, pickleworm, melonworm, downy mildew, and anthracnose. Apply 20 pounds of dust per acre. Do not use on watermelons. **DO NOT APPLY SAME DAY OF HARVEST.**

BEANS — Mexican bean beetle, bean leaf beetle, flea beetle, leafhoppers, tarnished plant bug and velvetbean caterpillar. Apply 40 pounds of dust per acre. **DO NOT APPLY SAME DAY OF HARVEST.**

CORN — Flea beetles, leafhoppers, corn rootworm adults (cucumber beetles) sap beetles, Japanese beetles, corn ear worm and European corn borer. Apply at rate of 40 pounds per acre. **DO NOT APPLY SAME DAY OF HARVEST; DO NOT FEED FORAGE TO DAIRY ANIMALS OR ANIMALS BEING FURNISHED FOR SLAUGHTER.**

APPLICATION OF THIS PRODUCT TO THE TASSEL REGION OF CORN DURING THE POLLEN SHED PERIOD WILL SERIOUSLY REDUCE BEE POPULATION.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.

SEVIN® — Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

NET WEIGHT

50 POUNDS

FC-898 A1271

PROD. 1540

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601



SEVIN ZINEB 3 - 5.2 DUST

INGREDIENTS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	3.0%
Zineb (zinc ethylene bisdithiocarbamate)	5.2%
INERT INGREDIENTS	91.8%
Total	100.0%

SEVIN—Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

CAUTION

Keep Out Of Reach Of Children

See Remainder of CAUTION Statement on Back Panel

USDA Reg. No. 1386-511

NET WEIGHT

4 POUNDS

FC-914-D967

Prod. 1543

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE OHIO

44-11-1969