

SEVIN[®] ZINEB
3-5.2 Dust

ACCEPTED
FEB 11 1983
Director for the Federal Insecticide,
Fungicide, and Rodenticide
Registration
EPA Reg. No. 1386-511


KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate)..... 3.0%
Zineb (zinc ethylene bisdithiocarbamate)..... 5.2%
INERT INGREDIENTS 91.8%
Total 100.0%

SEVIN - Trademark of Union Carbide Corp. for 1-naphthyl
N-methylcarbamate.

Net Weight: 40 Pounds
Product 1543

EPA Reg. No. 1386-511
EPA Est. No. _____



UNIVERSAL COOPERATIVES, INC.
MINNEAPOLIS, MN 55420

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Do not contaminate feed and foodstuffs. Protective clothing (long pants, long sleeved shirts, gloves, hat, and boots) must be worn during loading. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using; change clothing.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
If inhaled: Remove victim to fresh air. Apply respiration if indicated.
If on skin: Wash with soap and water. If in eyes: Flush eyes for at least 15 minutes with water. Call a physician.

NOTE TO PHYSICIAN

Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift of spray from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To control the insects and the plant diseases on the field-grown crops listed below, apply at the rate specified for each crop, to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough, but do not apply excessive amounts. Begin applications when the insects or their damage first appear and before diseases appear (when disease is first reported in the area). Repeat as necessary, generally at 7-day intervals unless a shorter interval is stated. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES - Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, early blight, late blight, leaf spot, and anthracnose on field grown tomatoes. Apply 40 to 50 pounds of dust per acre.

BEETS - Flea beetles, spittlebug, tarnished plant bug, stink bug, cabbage looper (when larvae are small, apply at 4 to 5-day intervals), and leaf spot. Apply 30 to 40 pounds of dust per acre. DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED; IF TOPS WILL NOT BE SO USED, DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

EGGPLANT AND PEPPERS - Leafhoppers, fall armyworm, European corn borer, tarnished plant bug, leaf spot, and phomopsis. Apply 30 to 40 pounds of dust per acre.

BEANS - Mexican bean beetle, bean leaf beetle, leafhoppers, rust, anthracnose and downy mildew (on lima beans). Apply 30 to 40 pounds of dust per acre. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

CABBAGE, CAULIFLOWER, AND BROCCOLI - Harlequin bug, imported cabbage worm, larvae of diamond-back moth, cabbage looper (when larvae are small; apply at 4 to 5-day intervals), and downy mildew. Apply 30 to 40 pounds of dust per acre. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

CUCUMBERS, PUMPKINS, AND SQUASH - Pickle worm, melon worm, cucumber beetles, flea beetles, leafhoppers, squash bug, downy mildew, *Alternaria* (*Macrosporium*) blight (on cucumbers), and anthracnose (on squash). Apply 30 to 33 pounds of dust per acre.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to approved Federal, State or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions, or warnings stated on this label.