

UNICO
MZ 5-5.2 DUST

1386-512
1.1.1.1

Contains 5% Methoxychlor and 5.2% Zineb

INGREDIENTS

ACTIVE INGREDIENTS:	
Methoxychlor, technical*	5.0%
Zineb (zinc ethylene bisdithiocarbamate)	5.2%
INERT INGREDIENTS	<u>89.8%</u>
Total	100.0%

*Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane
and 0.6% other isomers and reaction products

CAUTION (18-point type)

Keep Out of Reach of Children (12-point type)

See Remainder of CAUTION Statement on Back Panel

USDA Reg. No. 1386-512

Net Weight

5 Pounds

Prod 1542

Distributed By
UNITED CO-OPERATIVES, INC.
Alliance, Ohio

UNICO MZ 5-5.2 DUST

CAUTION

HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF NOSE AND THROAT!

DO NOT BREATHE DUST. AVOID CONTACT WITH SKIN. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land located well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

BEANS -- To control Mexican bean beetle, bean leaf beetle, leafhoppers, downy mildew, rust, and anthracnose, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear, and repeat at weekly intervals as necessary. Under severe disease conditions, applications at 4 to 5-day intervals may be necessary.

DO NOT APPLY TO BEANS WITHIN 7 DAYS OF HARVEST.

CABBAGE, CAULIFLOWER, AND BROCCOLI -- To control imported cabbage worm, Harlequin bug, downy mildew, and anthracnose, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover

upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear, and repeat at weekly intervals or more often under severe disease conditions.

DO NOT APPLY TO CABBAGE AND CAULIFLOWER WITHIN 7 DAYS OF HARVEST.

DO NOT APPLY TO BROCCOLI WITHIN 14 DAYS OF HARVEST.

CUCUMBERS, MELONS, AND SQUASH (Field Grown only) -- to control downy mildew, anthracnose, *Macrosporium* blight, and cucumber beetles, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear, which is usually when the runners start to form, and repeat at 5 to 10-day intervals depending upon severity of disease conditions and insect infestation.

DO NOT APPLY TO CUCUMBERS, MELONS, AND SQUASH WITHIN 1 DAY OF HARVEST.

TOMATOES (Field Grown only) -- To control early blight, late blight, anthracnose, gray leaf spot, *Septoria* leaf spot, blister beetles, and flea beetles, apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre. Begin dusting before diseases appear or before insects cause noticeable damage, which is usually shortly after transplanting into the field, and repeat at 5 to 10-day intervals or oftener under severe late blight conditions.

DO NOT APPLY TO TOMATOES WITHIN 1 DAY OF HARVEST.

NOTICE:

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.