

MZ 5-5.2 DUST
Contains 5% Methoxychlor and 5.2% Zineb


KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENTS:	
Methoxychlor, technical*	5.0%
Zineb (zinc ethylene bisdithiocarbamate)	5.2%
INERT INGREDIENTS	<u>89.8%</u>
Total	100.0%

*Equivalent to 4.4% 2,2-bis (p-methoxyphenol)-1,1,1-trichloroethane and 0.6% other isomers and reaction products.

NET WEIGHT: 40 POUNDS
PRODUCT 1542

EPA Reg. No. 1386-512
EPA Est. No. 1386-OH-1



UNIVERSAL COOPERATIVES, INC.
MINNEAPOLIS, MN 55420

ACCEPTED
1386-512

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or inhaled. Do not breathe dust. Avoid contact with eyes, skin, or clothing. Do not contaminate feed and foodstuffs. Protective clothing (long pants, long sleeve shirts, gloves, hats, and boots) must be worn during loading.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. If inhaled: Remove victim to fresh air. Apply respiration if indicated. If on skin: Wash with soap and water. If in eyes: Flush eyes for at least 15 minutes. Call a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift of dust from area treated. Do not apply where runoff may occur as a result of heavy rain. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEANS - To control Mexican bean beetle, bean leaf beetle, leafhoppers, downy mildew, rust, and anthracnose, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear, and repeat at weekly intervals as necessary. Under severe disease conditions, applications at 4 to 5-day intervals may be necessary.

DO NOT APPLY TO BEANS WITHIN 7 DAYS OF HARVEST.

CABBAGE, CAULIFLOWER, AND BROCCOLI - To control imported cabbage worm, Harlequin bug and downy mildew, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear, and repeat at weekly intervals or more often under severe disease conditions.

DO NOT APPLY TO CABBAGE AND CAULIFLOWER WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO BROCCOLI WITHIN 14 DAYS OF HARVEST.

CUCUMBERS, MELONS, AND SQUASH (Field Grown Only) - To control downy mildew, anthracnose, Alternaria (Macrosporium) blight, and cucumber beetles, uniformly apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre to cover upper and lower surfaces of foliage thoroughly. Begin applications when disease is first reported in area or when insects first appear which is usually when the runners start to form, and repeat at 5 to 10-day intervals dependent upon severity of disease conditions and insect infestation.

DO NOT APPLY TO CUCUMBERS, MELONS, AND SQUASH WITHIN 1 DAY OF HARVEST.

TOMATOES (Field Grown Only) - To control early blight, late blight, anthracnose, gray leaf spot, Septoria leaf spot, blister beetles, and flea beetles, apply 30 to 35 pounds of dust (1.50 to 1.65 lb. methoxychlor and 1.56 to 1.82 lb. zineb) per acre. Begin dusting before diseases appear or before insects cause noticeable damage, which is usually shortly after transplanting into the field, and repeat at 5 to 10-day intervals or oftener under severe late blight conditions.

DO NOT APPLY TO TOMATOES WITHIN 1 DAY OF HARVEST.

POTATOES - To control blister beetles, leaf hoppers, early blight, and late blight, apply 35 to 40 pounds of dust (1.65 to 2.0 lb. methoxychlor and 1.82 to 2.08 lb. zineb) per acre to foliage. Begin dusting when disease is first reported in area or when insects first appear, which is usually when potato plants are 2 to 6 inches high, and repeat at 5 to 10-day intervals depending upon severity of disease conditions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.