severe injury to hearby plants. Coarse sprays are less likely to drift. Do not apply to the soil in areas where the roots of desirable trees or ornamental plants may extend, or in locations where the chemical may be washed into contact with their roots. Do not apply within the drip line of trees and shrubs. Do not clean application equipment in these areas.

Do not graze or feed or sell for feeding to livestock and poultry any foliage from treated areas. Keep children and pets off treated areas until the herbicide has been washed into the soil by adequate rainfall and weed foliage has completely dried. If tall vegetation is treated, remove when dead to reduce hazard from fire.

Application equipment and utensils used to apply this herbicide should not be used subsequently for applying insecticides, fungicides, selective herbicides, and tertilizers, since it is almost impossible to clean such equipment adequately to prevent injury to desirable plants. Thoroughly clean and dry mechanical equipment after each use, however, to reduce corrusion.

CAUTION

Harmful if swallowed or absorbed through skin. Causes skin critation. Do not preathe vapor or spray mist. Do not get in eyes, on skin, or on clothing. In case of skin contact, wash with scap and water. If n eyes, flush with plenty of water and get medical attention. Wash thoroughly after handling. Avoid contamination of feed and toodstuffs. Do not store or use near heat or open flame. Do not store with or near feed, seeds, fertilizers, insecticides, and fungicides.

This product is poisonous to fish and wildlife. Keep out of any body of water Apply only for the purpose and in the manner stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions and unused chemical in a pit on popproductive splil located well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. It container is glass threak into a previously prepared put in an isolated area and buty tragments deeply. It container is metal, pinch holes in top and bottom, and crust under wheels of a heavy vehicle or with an axis and hory devoly in an isolated area will as in from water implies.

NOTICE

. Use only for the purpose and γ compliance with the finitations, a firm with right to this fabel.



KLEEN WALK Weed Killer

FOR NON-SELECTIVE CONTROL OF WEEDS IN AREAS WHERE VEGETATION GENERALLY MAY BE KILLED

ACTIVE INGREDIENTS

2-Methoxy 4,6 bis/isopropylamino,-sitriazine*

Pentachlorophenol .86%
Other chlorophenols and related compounds
Aromatic petroleum derivative solvent 91.00%
(NERT INGREDIENTS 6.55%)

Total 100.00
PROMETONE Herbicide TM Gelgy Chem. Corp., U.S. Par. 2 407-420

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel

USDA Reg No. 1386-524

Net Volume

. .-. !

1 U.S. Quart Liquid

PROD. 2105

H&G-124B-2-68

MANUFACTURED BY

UNICO KLEEN WALK WEED KILLER

DIRECTIONS FOR USE

USE THIS PRODUCT ONLY IN OR ON AREAS WHERE ALL PLANT GROWTH MAY BE PREVENTED FOR SEVERAL WEEKS TO AN ENTIRE GROWING SEASON. DO NOT USE ON OR NEAR DESIRABLE PLANTS OF CROPS, OR ON THE SOIL IN WHICH DESIRABLE PLANTS ARE GROWING OR ARE TO BE GROWN.

APP:ICATION To control most annual and shallow-rooted weeds and grasses, this product undiluted may be applied as a spray, or it may be mixed with water and the diluted mixture applied as a spray or with a sprinkling can.

The best time to apply is before weed seeds germinate or soon after weed emergence when the weeds are in a rapidly-growing succulent condition and are not over 2 to 4 inches tall. If unwanted vegetation is more than 4 inches high, cut or move to a height of about 2 inches, remove cut vegetation, and apply the spray to the remaining stubble and soil.

Attrough "burning" of plant to liage may be evident within a few days after application, this product acts primarily through the roots and therefore, sufficient rainfall to move the chemical into the root zone in the soil is necessary, and actual killing of weeds may require 2 to 3 weeks after adequate rainfall has occurred. Under conditions of frequently heavy rainfall or if the growing season is long a second application may be necessary to maintain seasonal control of weed.

RATES To control annual weeds and grasses on walks, driveways, patios, clay tennis rourts, and parking lots, apply 1 quart of undiluted product per 100 square feet $10^{\circ} \times 10^{\circ}$ of area to be treated. The spray should be applied uniformly and thoroughly to the area.

For firelanes, fence rows, and similar non-productive areas where vegetation is not dosired, dilute or mix 1 quart of product in 3 gallon of water and apply the total quantity of mixture to 100 square feet of area. The application may be made with a corayer or with a sprinkling can to effect uniform distribution and thorough coverage of the weed foliage and the soil. A second application may be required in 2 months to suppress deep rooted perennial weeds such as Canada thistle and milk must

For aport treatments on driveways, walks, and patios, apply the codiluted product is a seeds are noticed, as a coarse wetting spray.

EIV FATIONS Considerable caution must be exercised in using this material to avoid 10, may to crook and desirable plants. Keep the spray away from flower beds, tawns stribs, and other desirable vegetation. Apply only when there is no hazard from drift of spray from the area treated, since even minute quantities may cause

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO