

UNICO

5% METHOXYCHLOR DUST

INGREDIENTS

ACTIVE INGREDIENTS:

Technical methoxychlor *	5 %
INERT INGREDIENTS	95 %
Total	100 %

*Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related compounds

USDA Reg. No. 1386-

CAUTION

(18-Point Type)

KEEP OUT OF REACH OF CHILDREN

(12-Point Type)


Harmful if swallowed. Avoid breathing dust. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner directed in this label. Do not apply when weather conditions favor drift of dust from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of waste and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

BEANS -- To control Mexican bean beetle, corn earworm, and flea beetles, apply 30 to 40 pounds per acre. Begin applications before insects become numerous and repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO BEANS WITHIN 3 DAYS OF HARVEST. DO NOT FEED OR GRAZE TREATED BEAN PLANTS OR VINES TO LIVESTOCK.



4/23/68

1386-531

BEETS, CARROTS, EGGPLANT, LETTUCE, AND PEPPERS -- To control flea beetles, apply 30 to 35 pounds per acre to foliage. Begin applications when insects first appear and repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO BEETS AND CARROTS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED FOR FOOD OR FEED OR WITHIN 7 DAYS IF TOPS ARE NOT SO USED. DO NOT APPLY TO EGGPLANT AND PEPPERS WITHIN 1 DAY OF HARVEST. DO NOT APPLY TO LETTUCE WITHIN 14 DAYS OF HARVEST. *Comments*

CANTALOUPE, CUCUMBER, AND SQUASH -- To control cucumber beetles, apply 30 to 35 pounds per acre to foliage. Begin applications when insects first appear and repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO CANTALOUPE, CUCUMBER, AND SQUASH, WITHIN 1 DAY OF HARVEST.


PEA (GARDEN) -- To control alfalfa looper, celery looper, and pea weevil, apply 40 pounds per acre to foliage. Begin applications when insects first appear, and apply particularly during the early bloom stage if pea weevil is present. Repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO GARDEN PEAS WITHIN 7 DAYS OF HARVEST.

TOMATOES -- To control blister beetles, apply 30 to 35 pounds per acre to foliage when insects first appear. Repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO TOMATOES WITHIN 1 DAY OF HARVEST.

RASPBERRIES, BLACKBERRIES, BOYSENBERRIES, AND LOGANBERRIES -- To control Japanese beetle (adults), rose chafer, and strawberry weevil, apply 35 to 40 pounds per acre to foliage. Begin applications when insects first appear, and repeat at 7 to 10-day intervals as needed. DO NOT APPLY TO BLACKBERRIES, BOYSENBERRIES, LOGAN BERRIES, AND RASPBERRIES WITHIN 14 DAYS OF HARVEST.

STRAWBERRIES -- To control flea beetles, omnivorous leaf tier, spittlebug, strawberry leaf beetle and strawberry weevils, apply 30 to 35 pounds per acre of foliage. Begin applications when insects first appear and repeat at 7 to 10-day intervals as necessary. DO NOT APPLY TO STRAWBERRIES WITHIN 3 DAYS OF HARVEST.

-3-



NOTICE

Use only for the purposes and in compliance with the limitations, *Comments*
cautions or warnings stated on this label.

NET WEIGHT

50 Pounds
PROD 1093

Distributed By

UNITED CO-OPERATIVES, INC.
Alliance, Ohio