

UNICO

"Ready To Use"

WARFARIN

RODENTICIDE PELLETS

INGREDIENTS

ACTIVE INGREDIENTS:

WARFARIN [3-(a-acetylbenzyl)-
4-hydroxycumarin] 00.025%

INERT INGREDIENTS: 99.975%

Total 100.000%

CAUTION (18-point type)

Keep Out of Reach of Children (12-point type)

See Remainder of CAUTION Statement on Back Panel

USDA Reg. No. 1386-

Net Weight

5 Pounds
Prod. 904

Distributed By
UNITED CO-OPERATIVES, INC.
Alliance, Ohio

ACCEPTED
MAY 15 1968

UNICO Ready to Use Warfarin Rodenticide Pellets

5-15-60
1276-33

DIRECTIONS FOR USE

This product contains WARFARIN, an anticoagulant chemical, which reduces the clotting ability of the blood, and upon repeated feedings causes fatal hemorrhage in rats and mice.

RATS - Place 4 to 6 ounces of pellets in dry containers distributed throughout the areas or locations rats are known to frequent or are accustomed to feed. The number of containers (bait stations) needed depends upon the size of the area and the population of rats - use a sufficient number of stations next to burrows and in runways so the rats do not have to search for the pellets. Inspect bait stations daily, particularly during the first two weeks, and replenish the pellets as necessary to maintain an uninterrupted supply until all indications of feeding have stopped, usually a period of not less than 14 days.

MICE - Place 1 tablespoonful of pellets in shallow containers distributed at intervals of about every 6 to 8 feet in areas where mice or their signs have been seen. Inspect the pellets frequently and replenish as necessary in order to maintain an uninterrupted supply until all feeding has stopped, usually a period of 14 days.

SPECIAL NOTE - Always replace stale, damp, or moldy bait with fresh pellets. Burn or bury discarded bait.

If a source of continuous infestation of rats or mice prevails such as from nearby dumps or fields, permanent stations baited with the pellets should be used. Inspect the stations frequently and replace with pellets as needed.

CAUTION

Bait stations or pellet containers should be placed so that they are inaccessible to or protected from children, domestic animals, or pets. A protective device may be a short length of board firmly attached at an angle to a wall, beneath which the pellets are accessible to rats and mice but not to larger animals. Short

lengths of pipe or drain tile of suitable diameter also may be used.

Keep away from humans, domestic animals, and pets.

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose.

Burn empty containers (paper) in a hot fire, outdoors, in an isolated area.

The smoke is hazardous. Stay well away from the smoke.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

5-15-66
1300 533