

UNICO

Warfarin Rodenticide Concentrate

For Preparation of Mixed Baits to Kill Rats
and Mice


INGREDIENTS

ACTIVE INGREDIENTS:

WARFARIN [̢-(a-acetonylbenzyl)-4-hydroxycoumarin] . . . 0.5%

INERT INGREDIENTS 99.5%

Total 100.0%



CAUTION (18-point type)

KEEP OUT OF REACH OF CHILDREN (12-point type)

See Remainder of CAUTION Statement on Side Panel

USDA Reg. No. 1386-

NET WEIGHT

1 POUND
Prod. 902

Distributed By
United Co-Operatives, Inc.
Alliance, Ohio

UNION WARREN Rodenticide Concentrate

6/5/68

DIRECTIONS FOR USE

1386-534

This product is an anti-coagulant rat and mouse killer in concentrate form which should be mixed with a bait material suitable for rats and mice. It contains WARFARIN which reduces the clotting ability of the blood, and upon repeated feedings causes fatal hemorrhage in rats and mice.

MIXING BAITS - Mix thoroughly 1 part of the concentrate with 19 parts, by weight, of a bait material suitable for rats and mice. Most frequently used are corn meal, alfalfa meal, rolled oats, nut crumbs, poultry mash, or a mixture of these materials. A small amount, usually about 1 to 2 percent, of a bland oil such as mineral oil, corn oil, or other edible oil may be incorporated into the dry bait to reduce dustiness. Perishable materials, such as fresh ground meat, fish, cheese, and vegetables may be used, but baits made with these materials should be replaced daily. Dogs and cats will actually search out meat and fish. Baits made with perishable materials must be placed so as to positively prevent access by pets.

RATS - Place 6 to 8 ounces of mixed bait in dry containers distributed throughout the areas or locations rats are known to frequent or are accustomed to feed. The number of containers (bait stations) needed depends upon the size of the area, and the population of rats - use a sufficient number of stations next to burrows and in runways so the rats do not have to search for the bait. Inspect bait stations daily, particularly during the first week and replenish the bait as necessary to maintain an uninterrupted supply until all signs of feeding have stopped, usually a period of not less than 14 days.

MICE - Place 1 tablespoonful of mixed bait in shallow containers distributed at intervals of about every 6 feet in areas where mice or their signs have been reported. Inspect bait frequently and replenish as necessary until all signs of feeding have stopped.

SPECIAL NOTE - Promptly replace stale, damp, or moldy bait with fresh bait.
ALWAYS BURN OR BURY DISCARDED BAIT.

If a source of continuous infestation of rats or mice prevails such as from nearby dumps or fields, permanent stations baited with the dry bait should be used. Inspect the stations frequently and replace with bait as needed.

CAUTION

Bait stations or containers should be placed so that they are inaccessible to or protected from children, domestic animals, or pets. A protective device may be a short length of board firmly attached at an angle to a wall, beneath which the bait is accessible to rats and mice but not to larger animals. Short length of pipe or drain tile of suitable diameter also may be used.

Keep away from humans, domestic animals and pets.

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses by bishydroxycoumarin.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

6/5/68

1386-534