

ACCEPTED
 Apr 30, 1969
 UNDER NO. 1386-538

5% SEVIN 7 1/2% CAPTAN DUST

INGREDIENTS

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	7.5%
INERT INGREDIENTS	87.5%
Total	100.0%

USDA Reg. No. 1386-538

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not contaminate feed and foodstuffs.

NOTE FOR PHYSICIAN—Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner directed in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

For protection of honey bees, avoid application during flowering or pollinating period when bees may be visiting the crop. When necessary to apply during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after the application.

To control the insects and plant diseases on the crops listed below, apply at the rate stated in pounds per acre, uniformly and thoroughly to upper and lower surfaces of the foliage. Dust coverage of foliage should be complete, but not excessive. Begin applications before symptoms of diseases appear and when the insects or their damage first appear. Repeat as necessary, generally at 7-day intervals unless a shorter interval is required by the occurrence of weather conditions favorable to

from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit or non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire outdoors in an unpopulated area. The smoke is hazardous. Stay well away from the smoke.

For protection of honey bees, avoid application during flowering or pollinating period when bees may be visiting the crop. When necessary to apply during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after the application.

APPLICATIONS

To control the insects and plant diseases on the crops listed below, apply at the rate stated in pounds per acre, uniformly and thoroughly to upper and lower surfaces of the foliage. Dust coverage of foliage should be complete, but not excessive. Begin applications before symptoms of diseases appear and when the insects or their damage first appear. Repeat as necessary, generally at 7-day intervals unless a shorter interval is recommended by the occurrence of weather conditions favorable to the development of diseases.

GRAPES—Black rot, downy mildew, grape berry moth, Japanese beetle adults, and red banded leaf roller. Apply at rate of 40 to 60 pounds per acre in five or more applications timed as follows:

- (1) When buds or shoots show 1/2-inch green.
- (2) When shoots are 10 to 12 inches long.
- (3) One week after bloom.
- (4) Two weeks after bloom, and thereafter at 7 to 10 day intervals for one or more applications as needed.

STRAWBERRIES—Botrytis fruit rot, meadow spittle bug, strawberry leaf roller, and strawberry weevil. Apply at rate of 40 pounds per acre beginning as soon as new growth starts and repeat at approximately weekly intervals. Applications at intervals of 3 to 5 days may be required when frequent rains, heavy dews, or other conditions favorable to infection occur. Do not apply within 1 day of harvest.

CUCUMBERS, SQUASH, AND MELONS—Angular leaf spot, anthracnose, downy mildew, cucumber beetles, pickleworm, melonworm, leafhoppers, and squash bug. Apply at rate of 20 pounds per acre. Some leaf injury may occur if treatments are made when foliage is wet or during periods of prolonged high humidity. Do not use on watermelons.

POTATOES AND TOMATOES—Early blight, late blight, Colorado potato beetle, flea beetle, leafhoppers, fall armyworm, tomato fruitworm, tomato hornworm, and tarnished plant bug. Apply at rate of 40 pounds per acre beginning when potato plants appear above ground and when tomato plants are transplanted into the field. Repeat at approximately weekly intervals.

DURING PERIODS OF SEVERE INFECTION CONDITIONS, APPLICATIONS AT 3 TO 5 DAY INTERVALS MAY BE NECESSARY ON ALL THE CROPS ABOVE LISTED.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity, if several days duration following application.

NOTICE

Use only for the purposes and in the manner shown on the label. Do not use in violation of the label.

Net Weight

50 Pounds

NET WT.

NET WT.

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO