## **UNICO Dairy and Stock Fly Spray**

### DIRECTIONS FOR USE

UNICO Dairy and Stock Fly Spray contains CIODRIN (R) Insecticide and VA-PONA (R) Insecticide (DDVP) in a petroleum oil base. It is ready-to-use spray for application directly to lactating dairy cattle and beef cattle to control house flies. horn flies, stable flies, and mosquitoes, and to aid in control of face flies. Also may be used as a barn spray to control house flies.

DIRECT SPRAYING OF LIVESTOCK-Apply as a finely atomized mist uniformly distributed over each animal (dairy cattle and beef cattle) at a rate not to exceed 2 fluid ounces daily per animal. Use good spray equipment to spray each animal, preferably from the rear, directing the spray along both sides, including the belly, flanks, and withers, down each leg, over the back, and around the poll so that droplets are deposited on the face. Hold nozzle of sprayer 2 or 3 feet away from animal so as to deposit droplets on the hair. DO NOT WET THE HIDE. Repeat daily as needed. For milking dairy animals, apply the spray 20 to 30 minutes before beginning to mitk, or after milking has been completed. DO NOT SPRAY UDDERS OF LACTATING DAIRY ANIMALS.

BARN SPRAYING-To control house flies in dairy barns, loafing sheds, livestock barns, and stables, apply 2 fluid ounces per 1,000 cubic feet as a finely atomized mist uniformly distributed throughout the area. To reduce air movement as much as possible, it is preferable to close windows and doors before spraying; leave them closed for 10 minutes after spraying and then ventilate the area. Repeat daily, morning and afternoon, as needed. Do not apply as a barn spray if animals which have received direct application within 8 hours remain in the barn; remove such animals before spraying.

DO NOT USE IN HUMAN HABITATION. DO NOT CONTAMINATE FEED. DRINKING WATER, MILK, AND MILK HANDLING EQUIPMENT. DO NOT AP-PLY IN AREAS OR ROOMS WHERE MILK IS PROCESSED OR STORED.

DO NOT APPLY REGULARLY TO CALVES UNDER 6 MONTHS OF AGE. BRAHMAN CATTLE SHOULD NOT BE TREATED AS THEY MAY SHOW HYPER-SENSITIVITY TO ORGANIC PHOSPHATES.

# DAIRY AND STOCK FLY SPRAY

**ACTIVE INGREDIENTS:** 

Dimethyl phosphate of alpha-methylbenzyl	
3-hydroxy-cis-crotonate*	1.00%
3-hydroxy-cis-crotonate*2,2-Dichlorovinyl dimethyl phosphate**	
Related compounds**	
Petroleum hydrocarbons	
INERT INGREDIENTS	



\*CIODRINE Insecticide

\*\*Equivalent to 0.25% **VAPONA®** Insecticide

clothing. eyes get i and befor Harmiul Atropine i tion with

Avoid

DO NO

DESTI

If containe vehicle or land in ar on non-cre

9-9-70 1321-551

# CAUTION

KEEP OUT OF REACH OF CHILDREN See Remainder of CAUTION Statement on Side Panel USDA Reg. No. 1386-551

**Net Volume** 

FC-6H-8-70

**5 U.S. Gallons Liquid** MANUFACTURED BY

PROD. 301


100.00%

Use of warnings

# DAIRY AND STOCK FLY SPRAY

### **ACTIVE INGREDIENTS:**

Dimethyl phosphate of alpha-methylbenzyl	
3-hydroxy-cis-crotonate* 2,2-Dichlorovinyl dimethyl phosphate**	1.00%
2,2-Dichlorovinyi dimethyi phosphate**	
Petroleum hydrocarbons	
INERT INGREDIENTS	
<b>-</b>	100.000/



\*CIODRIN® Insecticide

\*\*Equivalent to 0.25% VAPONA® Insecticide

## CAUTION

KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel
USDA Reg. No. 1386-551

**Net Volume** 

5 U.S. Gallons Liquid

FC-6H-8-70

MANUFACTURED BY

PROD. 30:

100.00%

### CAUTION

Avoid inhalation of vapor and spray mist. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate water, feed, or foodstuffs. Harmiul if swallowed; induce vomiting immediately and get medical attention. Atropine is antidotal; 2-PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT USE, SPILL, POUR, OR STORE NEAR HEAT OR OPEN FLAME.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is metal punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe and bury crushed container deeply in a pit on non-crop land in an isolated area. If container is glass, break into a previously prepared pit on non-crop land in an isolated area, and bury fragments leeply.

9-9-70

### NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.