



1386-554



"51" DUST COPPER--ROTENONE

INGREDIENTS

ACTIVE INGREDIENTS:

Copper, expressed as metallic (in copper oxide)	5%
Rotenone	1%
Other cube resins	2%
INERT INGREDIENTS	92%

Total 100%

USDA Reg. No. 1386-554

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause irritation of nose and throat. Avoid inhalation of dust.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner directed in this label. Do not apply when weather conditions favor drift of dust from area to be treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty paper container in a hot fire, outdoors in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

CUCUMBERS, MELONS, SQUASH, AND PUMPKIN—To control cucumber beetles, downy mildew, anthracnose, scab, and bacterial wilt, apply 30 to 40 pounds of dust per acre. Begin applications when runners start to form or when disease is first reported in area, and repeat at 5 to 10-day intervals during growing season.

TOMATOES—To control early blight, late blight, gray leaf spot, Septoria leaf spot, and flea beetles, apply 30 to 50 pounds of dust per acre. Begin applications before diseases are expected to appear, or when insects first appear, usually shortly after transplanting into the field, and repeat at 5 to 10-day intervals during growing season.

POTATOES—To control flea beetles, early blight, and late blight, apply 35 to 50 pounds of dust per acre. Begin applications when insects first appear or before disease appear (when disease is first reported in the area), and repeat at 5 to 10-day intervals during growing season.

BEANS—To control bean leaf beetle and Mexican bean beetle, use 25 to 40 pounds per acre. Repeat at 7 to 12-day intervals as necessary.

CABBAGE—To control cabbage worms apply at rate of 25 to 40 pounds per acre. Repeat at 7 to 12-day intervals.

EGG-PLANT—To control Colorado potato beetle, flea beetles, alternaria, Phomopsis, leaf spot, anthracnose, use at the rate of 20 to 40 pounds per acre. Dust as soon as seedlings are up. Repeat at 7 to 14-day intervals as necessary.

PEPPERS—To control Colorado potato beetles, flea beetle, anthracnose, cercospora use at the rate of 20 to 40 pounds per acre. Dust as soon as seedlings are up. Repeat at 7 to 14-day intervals as necessary.

In all cases, the dust should be applied thoroughly and uniformly with particular attention to coverage on the under side of all foliage. **AT HIGH DOSAGE LEVELS PLANT FOLIAGE MAY SHOW SOME DAMAGE FROM COPPER.**

DO NOT APPLY TO THE ABOVE NAMED CROPS WITHIN 1 DAY OF HARVEST.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

NET WEIGHT

FC-968

50 POUNDS

Prod. 1972

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601