



KILL FLY DV

Contains Dipterex* and DDVP

INGREDIENTS

ACTIVE INGREDIENTS:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate*	0.5 %
2,2-Dichlorovinyl dimethyl phosphate	0.46%
Related compounds	0.04%

INERT INGREDIENTS

99.00%

Total 100.00%

*Reg. U.S. Pat. Off. by Farbenfabriken Bayer A.G.,

Chemagro Corp. Licensee, U.S. Pat. 2,701,225

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Back Panel

USDA Reg. No. 1386-555

NET WEIGHT

H&G-134-D471

5 POUNDS

PROD. 2185

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

UNICO KILL FLY DV

DIRECTIONS FOR USE

HOUSEFLIES-To control flies in or around DAIRY BARNs, MILK ROOMS, FOOD PROCESSING PLANTS, PATIOS, PICNIC AREAS, POULTRY HOUSES, STABLES, STOCK PENS, KENNELS, ANIMAL HOSPITALS, GARBAGE AREAS, AND OUTDOOR LATRINES, the bait should be used either as a dry bait ("as is") or as a liquid bait (dissolved in water).

DRY BAIT: Scatter lightly and uniformly in strips on floors, window sills, ledges, and other horizontal surfaces or areas. Use 4 ounces per 1,000 square feet (1 ounce per 250 square feet) distributed so that bait is barely visible over surfaces where flies congregate. Apply to surfaces of garbage and manure piles. Killing action may be enhanced if applied to moistened surfaces.

LIQUID BAIT: Mix 1 pound of bait in 1/2 gallon of water, add 1 cupful of corn syrup or 1/2 pound of sugar, and stir until completely dissolved. This makes an approximately 0.1% Dipterex bait. Sprinkle this water solution on floors, ledges, and horizontal surfaces as directed for dry bait, and on outside of farm buildings where fly population is intense or bothersome.

When floor area is lacking or is inadequate to assure uniform thorough distribution of the bait (dry or liquid) throughout the area (for example, a dirt-floor barn), the bait may be applied to previously moistened sacks, boards or strips of cardboard placed at all locations where flies congregate inside and outside the buildings. Vertical surfaces inside the buildings should be treated by spraying or painting liberally with the liquid bait. Use enough bait to assure uniform distribution in the area.

Maintenance of cleanliness and sanitation procedures, or good housekeeping practices, is essential to fly control in any or all situations. Promptly remove and dispose of manure and wastes in which flies breed. Apply the bait daily until flies are controlled and then repeat applications frequently as needed.

COCKROACHES-To control roaches in STORES, RESTAURANTS, AND SIMILAR COMMERCIAL BUILDINGS, place small quantities of the dry bait on pieces of paper, cardboard, or in shallow jar lids and distribute these at frequent intervals in secluded or protected areas where cockroaches hide or run, such as in cabinets or closets, under sinks, refrigerators, and counters. Each placement of the bait must be inaccessible to children and pet animals, and made so that dying insects cannot fall into food, utensils, or drinking water. Wet the bait lightly for enhanced killing action. Observe the bait placements frequently, and renew or replace as necessary.

LIMITATIONS

Do not apply directly to animals. Do not contaminate drinking water, milk, milk handling equipment and utensils. Do not treat portions of buildings which animals may lick. Do not apply to areas on or over which poultry feed. Do not feed treated garbage to animals. Applications directly to walls, floors, or other surfaces previously treated with lime, whitewash, or other alkaline materials may be ineffective. Keep bait and solution out of reach of domestic animals. Do not use in edible products areas of food processing plants.

CAUTION

Harmful if swallowed. Avoid contact with skin. Wash thoroughly with soap and warm water after using. Wash contaminated clothing with soapy and hot water before re-use. Do not contaminate feed and foodstuffs. If illness occurs get prompt medical aid. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE


Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

H&G-134-0471

PROD. 2185

Jun-25-1971

1386-555



KILL FLY DV

Contains Dipterex* and DDVP

INGREDIENTS

ACTIVE INGREDIENTS:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate*	0.5 %
2,2-Dichlorovinyl dimethyl phosphate	0.46%
Related compounds	0.04%

INERT INGREDIENTS	99.00%
Total	100.00%

Jun-25-1971

*Reg. U.S. Pat. Off. by Farbenfabriken Bayer A.G.,
Chemagro Corp. Licensee, U.S. Pat. 2,701,225

1386-555

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Back Panel

USDA Reg. No. 1386-555

NET WEIGHT

H&G-134-D471

5 POUNDS

PROD. 2185

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

UNICO KILL FLY DV

DIRECTIONS FOR USE

HOUSEFLIES To control flies in or around DAIRY BARNs, MILK ROOMS, FOOD PROCESSING PLANTS, PATIOS, PICNIC AREAS, POULTRY HOUSES, STABLES, STOCK PENS, KENNELS, ANIMAL HOSPITALS, GARBAGE AREAS, AND OUTDOOR LATRINES, the bait should be used either as a dry bait ("as is") or as a liquid bait (dissolved in water).

DRY BAIT: Scatter lightly and uniformly in strips on floors, window sills, ledges, and other horizontal surfaces or areas. Use 4 ounces per 1,000 square feet (1 ounce per 250 square feet), distributed so that bait is barely visible over surfaces where flies congregate. Apply to surfaces of garbage and manure piles. Killing action may be enhanced if applied to moistened surfaces.

LIQUID BAIT: Mix 1 pound of bait in 1/2 gallon of water, add 1 cupful of corn syrup or 1/2 pound of sugar, and stir until completely dissolved. This makes an approximately 0.1% Dipterex bait. Sprinkle this water solution on floors, ledges, and horizontal surfaces as directed for dry bait, and on outside of farm buildings where fly population is intense or bothersome.

When floor area is lacking or is inadequate to assure uniform thorough distribution of the bait (dry or liquid, throughout the area (for example, a dirt-floor barn), the bait may be applied to previously moistened sacks, boards or strips of cardboard placed at all locations where flies congregate inside and outside the buildings. Vertical surfaces inside the buildings should be treated by spraying or painting liberally with the liquid bait. Use enough bait to assure uniform distribution in the area.

Maintenance of cleanliness and sanitation procedures, or good housekeeping practices is essential to fly control in any or all situations. Promptly remove and dispose of manure and wastes in which flies breed. Apply the bait daily until flies are controlled and then repeat applications frequently as needed.

COCKROACHES To control roaches in STORES, RESTAURANTS, AND SIMILAR COMMERCIAL BUILDINGS, place small quantities of the dry bait on pieces of paper, cardboard, or in shallow jar lids and distribute these at frequent intervals in secluded or protected areas where cockroaches hide or run, such as in cabinets or closets, under sinks, refrigerators, and counters. Each placement of the bait must be inaccessible to children and pet animals, and made so that dying insects cannot fall into food, utensils, or drinking water. Wet the bait lightly for enhanced killing action. Observe the bait placements frequently, and renew or replace as necessary.

LIMITATIONS

Do not apply directly to animals. Do not contaminate drinking water, milk, milk handling equipment and utensils. Do not treat portions of buildings which animals may lick. Do not apply to areas on or over which poultry feed. Do not feed treated garbage to animals. Applications directly to walls, floors, or other surfaces previously treated with lime, whitewash, or other alkaline materials may be ineffective. Keep bait and solution out of reach of domestic animals. Do not use in edible products areas of food processing plants.

CAUTION

Harmful if swallowed. Avoid contact with skin. Wash thoroughly with soap and warm water after using. Wash contaminated clothing with soapy and hot water before re-use. Do not contaminate feed and foodstuffs. If illness occurs get prompt medical aid. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DESTROY EMPTY CONTAINERS: Do not reuse this container for any purpose. Burn empty containers in a hot fire outdoors in a well-ventilated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

For any further information, contact your supplier with the limitations of this product as stated on this label.

H&G 134-0471

PROD. 2185