



BAN AMINE WEED KILLER POST EMERGENCE

CORN (FIELD)—For the control of "phenoxy resistant" weeds such as, field bindweed, Canada Thistle, perennial sow thistle, horse nettle, prostrate spurge, smart weed, common milkweed, hedge bindweed, and other broadleaf weeds. For broadcast treatment—Use at the rate of 0.8 to 1.6 pints of product per acre, may be applied as a post emergence treatment over the tops of the corn plant (overall) until the corn is 10 inches high and then use drop nozzles and direct spray at base of corn plant. Do not apply after corn is 36 inches high and make one application per season. Use 7 to 25 gallons of water per acre. For Band Treatment (overall or directed), (1/3 of area) use at the rate of 0.26 to 0.52 pint (0.66 to 1.32 oz. dicamba acid equivalent plus 1.32 to 2.64 oz. 2,4-D acid equivalent).

One gallon of Ban Amine will treat 5 to 10 acres broadcast or 15 to 30 acres as a band treatment.

DO NOT USE ON SWEET OR POP CORN.

DO NOT GRAZE TREATED FIELDS OR HARVEST FOR DAIRY FEED PRIOR TO EN-SILAGE STAGE (MILK STAGE).

Fall Seeded Wheat (not undersowed to legumes)

For the control of "hard to kill" weeds (phenoxy resistant) including wild buckwheat, ladythumb, smartweed, knawel, corn cockle, mayweed, corn chamomile.

And for the control of other broadleaf weeds including Canada thistle (top growth), lambsquarters, pigweed, cocklebur, dragonhead (mint), mustards (annual), Rock, hempnettle, wild radish, wild garlic (top growth), vetch, Russian thistle, ragweed, stinkweed, goatsbeard (salsify), dandelion, Frenchweed (field pennycress), parakevine, field peppergrass (field pepperweed), sunflower and others.

Apply 0.8 pint Ban Amine (1/2 lb. dicamba plus 1/4 lb. 2,4-D) per acre. One gallon treats ten (10) acres.

Apply as a foliage spray immediately after winter dormancy and before the grain begins to joint.

Apply when weeds are actively growing. Apply in 32 to 40 gallons of water per acre.

LAWNS AND ORNAMENTAL TURF (not pasture)—To control chickweeds (common, mouse-ear and perennial or stitchwort), sheep sorrel, knotweed, clover, knawel, curly dock, dog fennel, chickory, mustard, ragweed, lambsquarters, plantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, black medick and stinging nettle in established perennial grasses in lawns or ornamental turf, use 1 fluid ounce (2 tablespoons) in 1 gallon of water and apply this mixture as a spray to wet uniformly the weed plants on 1200 square feet (40'x30'). This rate of application is equivalent to 2 1/4 pints per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition.

BAN AMINE WEED KILLER

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)* 15.2
Dimethylamine salt of related acids 2.8
Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid** 30.0

INERT INGREDIENTS 51.0

Total 100.0

*Equivalent to 12.69% 3,6-dichloro-o-anisic acid (dicamba 1.25#/gal.)

**Equivalent to 25.38% 2,4-dichlorophenoxyacetic acid (2.5#/gal.)

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional CAUTIONS

EPA Reg. No. 1386-559

Net Volume

5 U.S. Gallons Liquid

FC-872B-2-78

PROD. 15

MANUFACTURED FOR

UNIVERSAL COOPERATIVES, INC., ALLIANCE, OHIO 44601



BAN AMINE WEED KILLER

INGREDIENTS	
Methylamine salt of dicamba (3,6-dichloro-o-anisic acid)*	15.3%
Methylamine salt of related acids	2.4%
Methylamine salt of 2,4-Dichlorophenoxyacetic Acid**	30.5%
INGREDIENTS	51.8%
Total	100.0%

Equivalent to 12.65% 3,6-dichloro-o-anisic acid (dicamba 1.25#/gal.)
 Equivalent to 25.38% 2,4-dichlorophenoxyacetic acid (2.5#/gal.)

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional CAUTIONS

EPA Reg. No. 1386-559

Volume **5 U.S. Gallons Liquid**

PROD. 137

MANUFACTURED FOR

VERSAL COOPERATIVES, INC. ALLIANCE, OHIO 44601

Do not apply on or near desirable trees, shrubs, and other plants, or in locations where their roots may extend, or in locations where the chemicals may be washed or moved into contact with their roots. Do not clean application equipment or utensils in these areas. Do not apply to centipede grass, St. Augustine grass, or to Dichondra. Do not use on lawns where desirable bent grasses or clovers are present.

DO NOT GRAZE TREATED AREAS OR FEED TREATED FORAGE OR THRESHINGS TO LIVESTOCK. DO NOT APPLY TO NEWLY SEEDED LAWNS OR TURF.

CAUTION

HARMFUL IF SWALLOWED. DO NOT TAKE INTERNALLY. AVOID CONTACT WITH SKIN, EYES, AND CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES GET MEDICAL ATTENTION. DO NOT STORE WITH OR NEAR FERTILIZERS, SEEDS, INSECTICIDES, OR FUNGICIDES. Do not use or store near heat or open flame.

Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of this product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigating water supplies. Do not apply within the drip line of trees or shrubs.


It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e.g. wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops. Destroy empty containers; never re-use for any purpose.

This product is poisonous to fish. Keep out of any body of water. Apply only for the purposes and in the manner stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies. Do not burn empty containers.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.



desirable plants which are sensitive to spray mist to drift onto them since evaporation causes injury. Coarse sprays are less likely to cause injury. Vapors from this product after application in the immediate vicinity. Do not apply in a manner of producing puddles of solution in the soil. Excess of 2,4-D or silvex in the soil may temporarily retard plant growth. Do not contaminate domestic

~~This product is poisonous to~~

Apply only in the manner and for the purpose intended. Do not apply when weather conditions favor drift. Do not use where runoff may occur as a result of application. Clean equipment and dispose of rinse solution in a pit on non-crop land located well away from crops.

DESTROY EMPTY CONTAINERS. Do not reuse for any purpose. If container is glass, break in a prepared pit on non-crop land in an isolated area. If container is metal, punch holes in the bottom with the under wheels of heavy vehicle or with a sharp tool and bury deeply in a pit on non-crop land located well away from crops. Do not burn empty containers.

NOTICE

Use only for the purpose and in the manner and cautions or warnings stated on this label.



BAN AMINE WEED KILLER POST EMERGENCE

CORN (FIELD)—For the control of "phenoxy resistant" weeds such as, field bindweed, Canada Thistle, perennial sow thistle, horse nettle, prostrate spurge, smart weed, common milkweed, hedge bindweed, and other broadleaf weeds. For broadcast treatment—Use at the rate of 0.8 to 1.6 pints of product per acre, may be applied as a post emergence treatment over the tops of the corn plant (overall) until the corn is 10 inches high and then use drop nozzles and direct spray at base of corn plant. Do not apply after corn is 36 inches high and make one application per season. Use 7 to 25 gallons of water per acre. For Band Treatment (overall or directed), (1/3 of area) use at the rate of 0.26 to 0.52 pint (0.66 to 1.32 oz. dicamba acid equivalent plus 1.32 to 2.64 oz. 2,4-D acid equivalent).

One gallon of Ban Amine will treat 5 to 10 acres broadcast or 15 to 30 acres as a band treatment.

DO NOT USE ON SWEET OR POP CORN.

DO NOT GRAZE TREATED FIELDS OR HARVEST FOR DAIRY FEED PRIOR TO ENSILAGE STAGE (MILK STAGE).

Fall Seeded Wheat (not underseeded to legumes)

For the control of "hard to kill" weeds (phenoxy resistant) including wild buckwheat, ladysthumb, smartweed, knawel, corn cockle, mayweed, corn chamomile.

And for the control of other broadleaf weeds including Canada thistle (top growth), lambsquarters, pigweed, cocklebur, dragonhead (mint), mustards (annual), Ruchia, hempnettle, wild radish, wild garlic (top growth), vetch, Russian thistle, ragweed, stinkweed, goatsbeard (salsify), dandelion, Frenchweed (field pennycress), pasturevine, field peppergrass (field pepperweed), sunflower and others.

Apply 0.8 pint Ban Amine (1/2 lb. dicamba plus 1/4 lb. 2,4-D) per acre. One gallon treats ten (10 acres).

Apply as a foliage spray immediately after winter dormancy and before the grain begins to joint.

Apply when weeds are actively growing. Apply in 32 to 40 gallons of water per acre.

LAWNS AND ORNAMENTAL TURF (not pasture)—To control chickweeds (common, mouse-ear and perennial or stitchwort), sheep sorrel, knotweed, clover, knawel, curly dock, dog fennel, chickory, mustard, ragweed, lambsquarter, pliantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, Black medic, and stinging nettle in established perennial grasses in lawns or ornamental turf, use 1 fluid ounce (2 tablespoons) in 1 gallon of water and apply this mixture as a spray to wet uniformly the weed plants on 1200 square feet (40'x30'). This rate of application is equivalent to 2 1/4 pints per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition.

BAN AMINE WEED KILLER

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o anisic acid)*	15.3%
Dimethylamine salt of related acids	2.4%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid**	30.5%
INERT INGREDIENTS	51.8%
Total	100.0%

*Equivalent to 12.69%, 3,6-dichloro-o-anisic acid (dicamba 1.25#/gal.)

**Equivalent to 25.38%, 2,4-dichlorophenoxyacetic acid (2.5#/gal.)

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional CAUTIONS

EPA Reg. No. 1386-559

Net Volume

5 U.S. Gallons Liquid

FC-973B-2-78

PROD. 137

MANUFACTURED FOR

UNIVERSAL COOPERATIVES, INC. ALLIANCE, OHIO 44601

EPA Reg. No. 1386-559