

UI

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes and clothing. In case of contact, wash exposed skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not store near other pesticides or seed.

PESTICIDE DISPOSAL — Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL — Do not reuse empty container. The empty containers should be rinsed with water at least three times before disposal, and rinse water should be added to spray tank. Dispose of container in an approved landfill or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures.

NOTICE

Because Universal Cooperatives, Inc. has no control over storage, handling and conditions of use, which are of critical importance, Universal Cooperatives, Inc. makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Universal Cooperatives, Inc. assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.



BA

ACTIVE INGREDIENTS:
Dimethylamine salt of dk
Dimethylamine salt of rd
Dimethylamine salt of 2,
INERT INGREDIENTS

Total
Equivalent to 12.5%, 3,
Equivalent to 24.5%, 2
***ISOMER SPECIFIC BY**

ACCEPTED
13 MAR 12 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1386-559

KEEP OUT

See Side Panel

Net Contents

901373-1-81

78



UNIVE
MI

UNICO®

PRECAUTIONARY STATEMENTS HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes and clothing. In case of contact with skin with plenty of soap and water; flush eyes with water for 15 minutes and get medical attention. Keep out of

ENVIRONMENTAL HAZARDS

Do not contaminate water, streams or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not store in water, food or feed by storage, disposal or cleaning of equipment is prohibited. Do not store near other pesticides or

Pesticide, spray mixture or rinseate that cannot be reused should be disposed of in a landfill approved by local authorities in a safe place away from water supplies.

Do not reuse empty container. The empty container should be rinsed with water at least three times before disposal. Do not add to spray tank. Dispose of container in a safe place. Consult Federal, State or local authorities for approved alternative procedures.

NOTICE

Universal Cooperatives, Inc. has no control over storage, handling or use of this material. In an area of critical importance, Universal Cooperatives, Inc. makes no warranty, either express or implied, for results or damages which may be established by appropriate authorities. Universal Cooperatives, Inc. assumes any responsibility for damage to animals, soil or property arising out of misuse, abuse or improper handling of this material.

BAN AMINE WEED KILLER

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)*	15.06%
Dimethylamine salt of related acids	3.75%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid**	29.60%
INERT INGREDIENTS	51.59%

Total 100.00%

*Equivalent to 12.5%, 3,6-dichloro-o-anisic acid or 1.25 lbs./gal.
**Equivalent to 24.58%, 2,4-dichlorophenoxyacetic acid or 2.38 lbs./gal.
***ISOMER SPECIFIC BY AOAC METHOD NO. 6. DO1-5

ACCEPTED
MAR 12 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act registered under EPA Reg. No. 1336-559

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 1336-559
EPA Est. No. 1336-OH-1

Net Contents

78

5 U.S. Gallons Liquid (18.927 L)

901373-1-81

Manufactured By

PROD 137

UNIVERSAL COOPERATIVES, INC.
Minneapolis, Minnesota 55420

UNICO®

BAN AMINE WEED KILLER

CONTENTS:
salt of dicamba (3,6-dichloro-o-anisic acid)* 15.06%
salt of related acids 3.75%
salt of 2,4-Dichlorophenoxyacetic Acid** 29.80%
INERTS 51.59%
..... 100.00%
12.5% 3,6-dichloro-o-anisic acid or 1.25 lbs./gal.
24.5% 2,4-dichlorophenoxyacetic acid or 2.38 lbs./gal.
ANALYZED BY AOAC METHOD NO. 6. D01-5

OUT OF REACH OF CHILDREN

CAUTION

See Panel for Additional Precautionary Statements

EPA Reg. No. 1300-009
EPA Est. No. 1300-001-1

5 U.S. Gallons Liquid (18.927 L)

Manufactured By

K-2

PROD. 137

OPERATIVES, INC.

M... 5542

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

POST EMERGENCE

CORN (FIELD) (Ground Application) — For the control of "phenoxy resistant" weeds such as field bindweed, Canada thistle, perennial sow thistle, horse nettle, prostrate spurge, smartweed, common milkweed, hedge bindweed, and other broadleaf weeds. For Broadcast Treatment — Use at this rate of 0.8 to 1.8 pints of product per acre. May be applied as a post emergence treatment over the tops of the corn plant (overall) until the corn is 10 inches high and then use drop nozzles and direct spray at base of corn plant. Do not apply after corn is 30 inches high and make one application per season. Use 7 to 23 gallons of water per acre. For Band Treatment (overall or directed), (1/3 of area), use at the rate of 0.26 to 0.52 pint.

One gallon of Ban Amine will treat 5 to 10 acres broadcast or 15 to 30 acres as a band treatment.

DO NOT USE ON SWEET OR POP CORN.

Do not graze treated fields or harvest for dairy feed prior to ensilage stage (milk stage).

FALL SEEDED WHEAT (not underseeded to legumes) (Ground Application) — Do not graze or harvest for dairy feed prior to crop maturity.

For the control of "hard to kill" weeds (phenoxy resistant) including wild buckwheat, ladythumb, smartweed, knotweed, corn cockle, mayweed, corn chamomile.

And for the control of other broadleaf weeds including Canada thistle (top growth), lambsquarters, pigweed, cocklebur, dragonhead (mint), mustards (annual), Kochia, hempnettle, wild radish, wild garlic (top growth), vetch, Russian thistle, ragweed, stinkweed, goatsbeard (salsify), dandelion, Frenchweed (field pennycress), pasturevine, field peppergrass (field pepperweed), sunflower and others. Do not graze or harvest for dairy feed prior to crop maturity.

Apply 0.8 pint Ban Amine per acre. One gallon treats ten (10) acres.

Apply as a foliage spray immediately after winter dormancy and before the grain begins to joint.

Apply when weeds are actively growing. Apply 2 to 40 gallons of water per acre.

LAWNS AND ORNAMENTAL TURF (not pasture) — To control chickweeds (common, mouse-ear and perennial or stitchwort), sheep sorrel, knotweed, clover, knotweed, curly dock, dog fennel, chickory, mustard, ragweed, lambsquarters, plantain, daisy fleabane, pigweed, wild radish, bull thistle, burdock, dandelion, black medic, and stinging nettle in established perennial grasses in lawns or ornamental turf, use 1 fluid ounce (2 tablespoons) in 1 gallon of water and apply this mixture as a spray to wet, dormant turf plants on 1200 square feet (40' x 30'). This rate of application is equivalent to 1/4 pint per acre. Apply in the spring or early summer when weeds are in a succulent, rapidly growing condition. Do not apply to newly seeded lawns or turf.

Do not apply on or near desirable trees, shrubs, and other plants, or on areas where their roots may extend, or in locations where the chemicals may be washed or moved into contact with their roots. Do not clean application equipment or utensils in these areas. Do not apply to centipede grass, St. Augustine grass or to dichondra. Do not use on lawns where desirable bent grasses or clovers are present.

Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamental, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of this product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigation water supplies. Do not apply within the drip line of trees or shrubs.

It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from aer-solastic parts, e.g. wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops.

79