


1386 512



JUNIPER - juniper scale, crawlers only, spider mites
 LILAC - Powdery mildew
 PEONY - leaf spots, botrytis blight, phytophora blight, leafhoppers. For best results in control of phytophora keep foliage protected and liberally dust soil and crowns when top growth is removed, burn all plant residues
 PETUNIA - Japanese Beetle adults
 ROSE - Japanese Beetle adults, exposed thrips, leaf hopper, aphids, lacebug, powdery mildew, black spot.
 SNAPDRAGON - Powdery mildew, rust, plant bugs
 SPIREA - Aphids, leafhoppers, plant bugs
 ZINNIA - Japanese Beetle, alternaria leaf spot, powdery mildew, plant bugs

ROSE and FLORAL Dust No. 2



SPECIAL NOTES

Sulphur may cause injury to foliage if applied during periods of high daytime temperature. Do not apply during periods when the daytime temperature is 90 degrees F. or above, or within two days of the time such temperature may be expected.

Carbaryl Sevin is harmful to Boston Ivy. Keep dust away from Boston Ivy.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

CAUTION

Harmful if inhaled or swallowed. May cause irritation of eyes, nose, throat or lungs. Avoid contact with or breathing of dust. Wash thoroughly after handling and before eating or smoking. Take shower or bath after using, change clothing.

This product is poisonous to fish or wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies. Avoid contamination of feed and feedstuffs.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Destroy container in compliance with local regulations. Keep immediately away from smoke, smoke may be hazardous.

BEES - Honeybees should be treated at safe distance from all areas to be treated during the growing season. Avoid applications during periods when beneficial bees are visiting plants or blossoms.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

UNICO ROSE and FLORAL DUST NO. 2

DIRECTIONS FOR USE

Apply for application to ROSES, ORNAMENTAL TREES, SHRUBS and SHRUBS for control of the following plant diseases:

Apply during the early morning when plants are wet with dew or in the evening when the dew is gone. Begin applications before diseases are apparent and before insects become numerous, which is usually early in the season when new growth is just starting in the spring, when seedlings appear above ground, or when young plants are transplanted. Cover all foliage evenly and thoroughly with a light film of dust including underside of leaves. Repeat application at 5 to 10 day intervals and after heavy rains to maintain protection during the growing season unless directed otherwise for specific pests in following listing. To enhance disease control remove and destroy all prunings and plant residues by burning if permitted.

For application to the following plants and to control the following pests:

ASTER - leafhoppers, rust, botrytis, aster yellow, aster may be reduced by control of leafhoppers.

ARBORVITAE & TAXUS (YEW) - Bagworm, young in the larvae only, aphids, arborvitae leaf miner, juniper and Fletcher scale, crawlers, only, and mealy bug.

AZALEA, CAMELIA, RHODODENDRON - Rose of sharon, scale, aphids, lace bugs, exposed thrips, crawler stage of scales. For botrytis flower blight on azalea and rhododendron, dust flowers and liberally dust soil and crown of plants.

BOXWOOD, EUONYMUS - Leafminer, crawler stage of euonymus scale.

CHRYSANTHEMUM - leafhopper, aphid, spider mite, leaf spot, ray blight, plant bugs.

CLEMATIS - aster beetle, plant bugs.

DAHLIA - Japanese Beetle adults, leafhopper, mealy bug, mold, powdery mildew, plant bugs.

FLOWERING RABBIT - European spruce sawfly, European spruce sawfly and immediate control of the sawfly as soon as small webbing is started. For control of the sawfly protect until June 1st.

GLADIOLUS - European spruce sawfly, spider mite, botrytis blight.

HONEY LOCUST & MULBERRY - Mealy bug, crawler stage of mealy bug, scale, spider mite, leafhopper, aphid, keep foliage protected from insects during the growing months.

HOLLY - leafminer.
 IRIS - Alternaria leaf spot, spider mite.

INGREDIENTS

ACTIVE INGREDIENTS	
Carbaryl 1-naphthyl N-methylcarbamate	5.0%
Zineb zinc ethylene bisdithiocarbamate	5.0%
Sulphur	20.0%
INERT INGREDIENTS	
	70.0%
Total	100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1386-562

NET WEIGHT

1 POUND

H&G 139

Prod. 2155

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC.
 ALLIANCE, OHIO 44601