




1277-585

FRUIT SPRAY TREE - TRELLIS - GARDEN



GRAPES — Grape berry, both Japanese beetle adults and banded leaf roller, leathopper, two spotted spider mite, black rot, downy mildew.

Apply: 1. when buds or shoots show 1/2 inch green; 2. when shoot are 10 to 12 inches long; 3. one week after bloom; and 4. thereafter at 7 to 10 day intervals for one or more applications as needed. Do not apply within 7 days of harvest.

STRAWBERRIES — Spittlebug, strawberry leaf roller, strawberry weevil, two spotted mites, strawberry spotted mites, cyclamen mites, Botrytis fruit rot, and leaf spot.

Apply: 1. when new growth starts in the spring; 2. at subsequent 7 to 10 day intervals as needed. Sunrise and Early Dawn varieties may be injured by Sevin. Do not apply within 2 days of harvest.

RASPBERRY AND BLACKBERRY — Japanese beetle adults, leaf roller, leathoppers, raspberry aphid, European red mite, raspberry only, and raspberry anthracnose. Post harvest applications are necessary to control anthracnose. Attention to strict sanitation practices.

Apply: 1. when new growth starts; 2. at petal fall; at 7 to 10 day intervals. Do not apply within 7 days of harvest.

Do not graze or feed forage from treated areas to livestock.

SPECIAL NOTES

Not effective if used with alkaline materials. Avoid spraying when poor drying conditions are prevalent. Some injury may occur on tender foliage.

BEES — This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust or spray mist. Do not take internally. Avoid contact with skin. Wash before eating or smoking; take shower or bath after using; change clothing. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply only in the manner and for the purposes directed by this label. Do not apply when weather conditions favor drift of spray from areas treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS — Do not reuse this container for any purpose. Destroy container in compliance with local regulations. If burned stay away from smoke. Smoke may be hazardous.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions, or warnings stated on this label.

INGREDIENTS

ACTIVE INGREDIENTS	
Carbaryl 1-naphthyl N-methylcarbamate	17.5%
1,1-Bis p-chlorophenyl-2,2,2-trichloroethanol	6.0%
Captan N-trichloromethylthio 4-cyclohexene-1,2-dicarboximide	17.5%
INERT INGREDIENTS	
	59.0%
Total	100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE REMAINDER OF CAUTION STATEMENT ON SIDE PANEL

EPA Reg. No. 1386-565

Each Pound Makes 16 Gallon of Spray

NET WEIGHT 1 POUND
H&G 141 Prod. 2171

DISTRIBUTED BY

UNICO FRUIT SPRAY (TREE - TRELLIS - GARDEN)

DIRECTIONS FOR USE

Apply as a spray by mixing with water. Use 6 level, cable- spoonfuls per gallon of water (approx. equivalent to 6 lb. of product per 100 gal. of water). Application rate per acre will depend upon tree or plant size, age, plant vigor and row width. Mix thoroughly before beginning to spray, maintain agitation while spraying. Apply uniform and thoroughly to foliage and fruits. Proper timing of applications and uniformly thorough coverage are essential. Avoid spraying when poor drying conditions are prevalent; foliage injury may result.

The following insects, mites, and plant diseases can be controlled by proper applications as indicated:

APPLES AND PEARS — Codling moth, apple maggot, red banded leaf roller, green apple aphid, rosy apple aphid, European red mite, leathopper, two spotted spider mites, Japanese beetle adults, scab, Brooks fruit spot, sooty blotch, black rot, bitter rot, tent caterpillars, pear slug, budmoths and crawler stages of San Jose, oyster shell.

Apply: 1. delayed dormant; 2. just before bloom; when most of the petals have fallen; 3. 7 days after petal fall; and 4. at 7-10 day intervals as needed in cover sprays. Do not apply within 7 days of harvest.

Do not apply to D'Anjou pear, Base pear, Red D'Anjou, McIntosh, York or other varieties of apples and pears sensitive to Captan or Sevin may develop russet or show foliage injury from early sprays. Fruit thinning on apple varieties may occur from sprays applied during winter period following full bloom.

PEACHES, CHERRIES, APRICOTS, PLUMS — Oriental fruit moth, red banded leafroller, European red mite, two spotted mite, crawler stages of San Jose and oyster shell scales, brown rot, scab, leaf spot, Japanese beetle adults, peach twig borer and lesser peach tree borer, Peach tree borer. 4 sprays applied to limb bases and trunk areas in late summer, usually July and August, will provide protection from tree pests. Do not apply more than 500 gallons per acre on peaches, apricots, and plums, and 600 gallons per acre on cherries. Do not repeat application on plums within 30 days.

Apply: 1. when buds show pink; 2. just before buds open; 3. when most of the petals have fallen; 4. 10 to 15 days after petal fall; 5. on seven to ten day schedule as needed.

Do not apply within 14 days of harvest. Do not repeat applications on cherries within 30 days.

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601