

FOR THE GENERAL...  
1356 568



# LIVESTOCK PRESSURIZED SPRAY

**CONTAINS KORLAN\***  
**WOUND SPRAY**  
Kills Screw Worms—Blowfly Maggots  
FOR LIVESTOCK OF ALL AGES  
NEWBORN TOO

<b>ACTIVE INGREDIENTS:</b>	
Ronnel 0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate .....	2.5%
Xylene .....	.5%
<b>INERT INGREDIENTS:</b> .....	97.0%

\*Trademark of the Dow Chemical Company

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
Read Complete Precautions on Side Panel  
EPA Reg. No. 1386-568

**Net Weight 1 Pound (16 oz.)**

FC-989G-12-72      MANUFACTURED BY      PROD. 901

**UNIVERSAL COOPERATIVES, INC.**  
**ALLIANCE, OHIO 44601**

## LIVESTOCK REMEDY —

**NCT FOR HUMAN USE  
FOR EXTERNAL USE ONLY**

### DIRECTIONS FOR USE

For use on cattle, sheep and horses. Hold bomb about 3 inches from wound or infested area. Direct spout of container toward the area to be treated. Press valve down to release spray. Wet thoroughly by spraying directly into and about 1 inch around wound.

**SCREW WORM AND BLOWFLY INFESTATIONS:** Use particular care to get spray into deep pockets caused by screw worms. Repeat application 5 to 7 day intervals until the wound is healed.

**TO PREVENT SCREW WORM AND BLOWFLY INFESTATIONS:** Apply directly to all open or fresh wounds such as from dehorning, castration, shear cuts, etc. Treat navals of newborn animals. Freely bleeding wounds may require several treatments.

**FLEECEWORM INFESTATIONS:** Spray infested area liberally, working toward the center of the infestation. Shearing or removing the wool from the infested area is not necessary. Repeat every 3 days until infestation is controlled.

### CAUTION

**INJURIOUS TO EYES.** Do not get in eyes. Avoid breathing spray mists. Wash hands thoroughly after using. Avoid contamination of food and feedstuffs. Keep out of reach of children.

In case of human eye contact, flush eyes immediately with plenty of water and prompt medical attention.

Korlan is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is an antidote. Consult veterinarian at first sign of an adverse reaction.

### WARNING

Contents under pressure; do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

### NOTICE

Use only for the purposes and in compliance with limitations, cautions or warnings stated on this label.

