



CERTIFICATION OF COMPLIANCE

I, being an authorized representative of Universal Cooperatives, Inc., understand the labeling requirements specified in the Registration Standard for 2,4-Dichlorophenoxyacetic Acid as amended and included therein by reference. Further, I certify that all containers of Turf Treeter "T", EPA Reg. No. 1386-569, released for shipment after March 1, 1991, will bear revised labeling in accordance with those requirements.

UNIVERSAL COOPERATIVES, INC.
Attn: Ms. Nik Ramswick
Regulatory Affairs Technician
7801 Metro Parkway
P. O. Box 460
Minneapolis, Minnesota 55440


June 11, 1990

BEST AVAILABLE COPY



2074

TURF TREETER "T"

For "Hard-To-Kill"
Broadleaf Weeds
In Turf.



Active Ingredients:


*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	3.23%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	10.59%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) ...	1.28%
Inert Ingredients	84.90%
Total	100.00%

*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%. **2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76%. ***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Side Panel For Additional Cautions



BEST AVAILABLE COPY



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes, or clothing, avoid breathing spray mist. If contact with eyes occurs, flush with plenty of water.

Wear face shield or goggles, chemical resistant gloves, long pants, long-sleeved shirt, shoes and gloves when mixing, loading, or applying this product. Do not reenter or permit others to reenter treated areas until sprays have dried or dusts have settled.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Contact a physician immediately. Give victim one or two glasses of water and induce vomiting by touching the back of throat with a finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. **If On Skin:** Remove contaminated clothing and wash affected areas with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists. **If In Eyes:** Flush with water for at least 15 minutes. Call a physician immediately. **If Inhaled:** Remove victim to fresh air. Apply respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Keep out of lakes, streams, or ponds. Do not contaminate any body of water, especially domestic and irrigation water supplies.

Drift or run-off may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Use only lawn sprayers for this product. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPP and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

When cleaning equipment, do not pour the washwater on the ground, spray or drain over a large area away from wells and other water sources.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ONE QUART TREATS 6,400-12,800 SQUARE FEET OF TURF

Turf Treeter "T" is recommended for use on most turf grasses, especially those species sensitive to 2,4-D such as St. Augustine, bent, and buffalo. Temporary foliage discoloration of the preceding species may result from applications during periods of high temperatures. St. Augustine grass may show temporary discoloration from application of this product when soil moisture is excessively high. Discoloration might be more pronounced when higher dosages are applied either purposely or accidentally.

If soil moisture is low, water or irrigate turf areas 2 to 3 days prior to spraying with TURF TREETER "T". Do not water or irrigate for 24 hours after application is made. Spray on calm day to avoid drift, use a coarse wetting

spray when temperature is near 70° F. for best results. A larger volume of water will normally insure better coverage; however, water volume per unit of area is of less significance in control than using the correct amount of TURF TREETER "T" for the designated area. Weed species are more easily controlled when young and rapidly growing. More mature weeds will require higher dosage and repeat applications.

TURF TREETER "T" will injure or kill clover or dichondra lawns if spray is applied directly or permitted to drift on such areas. Spray drift or direct application will injure or kill ornamental and vegetable crops. Do not apply to new lawn seedlings until after seedlings are well established.

HOW TO MIX:

1. For use on St. Augustine Grass, bentgrass, buffalo grass, and grass seedlings mix 2 oz. in 1 gallon of water to uniformly spray 800 square feet; for example, an area 20 feet wide and 40 feet long. If necessary, repeat treatment in 2 to 3 weeks, if resistant weeds persist.

2. For use on other grasses such as bermuda, bahia, bluegrass, centipede, fescue, and rye mix 1 1/2 oz. in sufficient water to spray uniformly 400 square feet. On resistant species, under poor growing conditions, or for older weeds mix 2 oz. in sufficient water to spray 400 square feet with special attention to thorough coverage.

LIMITATIONS: Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter. Do not feed foliage or plant parts from treated areas to livestock.

WEEDS CONTROLLED—TURF TREETER "T" is especially effective against weeds such as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS, AND HENBIT. In addition, it kills most broadleaf weeds, some of which are listed below:

Bedstraw	Cocklebur	Lambsquarter	Poison Ivy	Thistle
Black Medic	Dandelion	Lespedeza	Poison Oak	Spurge
Buckhorn	Dock	Mallow	Purslane	Velvet Leaf
Bull Thistle	Galinsoga	Morning Glory	Ragweed	Wild Carrot
Burdock	Ground Ivy	Mustard	Sheep Sorrel	Wild Garlic
Buttercup	Hawkweed	Oxalis	Shepherd's	Wild Lettuce
Chicory	Heal-all	Peppergrass	Purse	Wild Onion
Chickweed	Henbit	Pigweed	Speedwell	Yarrow
Clover	Knotweed	Plantain	Tansy Mustard	

And many other broadleaf weeds.

STORAGE AND DISPOSAL

STORAGE: Store product in original container and in a locked storage area. Do not store with or near feed or food products, or in contact with fertilizer, seeds, insecticides, or fungicides. KEEP FROM FREEZING.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and **Buyer** assumes the risk of any use contrary to such directions. **Seller** makes no other express or implied warranty, including any other express or implied warranty of **Fitness** or of **Merchantability**, and no agent of **Seller** is authorized to do so except in writing with a specific reference to this warranty. In no event shall **Seller's** liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

EPA Reg. No. 1386-569
EPA Est. No. 1386-OH-1
Product 2103
Net Contents 1 Quart

Contains Trimec® a registered trademark of P.B.I.-Gordon Corporation, U.S. Patent No. 3,284,186
CO-OP is a registered trademark of Universal Cooperatives, Inc.

UNIVERSAL COOPERATIVES, INC., MINNEAPOLIS, MN 55440