

dew, Botrytis Grey Mold: Use 1 tablespoonful per 2 gallons of water. Begin application when disease first appears and repeat at 10 to 14 day intervals as long as needed.

TURF-Dollar Spot (Schlerotinia homeocarpo), Large Brown Patch (Rhizoctonia solani), Fusarium Patch (Fusarium Nivale): Apply at the rate of 2 oz. (12 tablespoonfuls), per 1,000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gallons per 1,000 sq. ft. Apply at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days. Do not graze or feed clippings from treated areas to livestock.

IMPORTANT

Never allow UNICO Systemic Rose & Ornamental Fungicide Spray to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of UNICO Systemic Rose & Ornamental Fungicide Spray as a fungicide. Keep container closed when not in use. Store in cool dry area.

CAUTION

Avoid breathing dust. Avoid contact with skin, eyes and clothing. May cause irritation of eyes, nose, throat and skin. Wash thoroughly after using. In case of contact; flush skin or eyes with plenty of water; for eyes get medical attention.

Keep away from fire or sparks.

ACCEPTED
DESTROY EMPTY CONTAINER. Do not reuse this container for any purpose. Break glass container into previously prepared pit in an isolated area well away from water supplies and bury fragments deeply.

Now NOTICE 12, 1972
Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1386-571

THIS PRODUCT FOR AGRICULTURAL USE.

U.S.D.A. Reg. No. 1388-23

DIRECTIONS: To prepare a Bordeaux Mixture use equal quantities of powdered Bluestone and Hydrated Spray Lime. Fill the tank one quarter full and with agitator running wash the Bluestone into the tank through a bronze or copper screen. Fill the tank three quarters full and wash the Hydrated Spray Lime through the screen and fill the tank. Agitate for several minutes to insure thorough mixing. A 5-5-50 Bordeaux Mixture (5 pounds Powdered Bluestone - 5 pounds Hydrated Spray Lime - 50 gallons water) is recommended for the control of Shot-Hole Fungus on Almonds, Apricots, Peaches, Nectarines, and for the control of Brown Rot on Almonds, Apricots, Cherries, Peaches, Nectarines, Plums and Prunes. For Shot-Hole Fungus apply as a dormant spray in late fall or early spring. For Brown Rot apply when buds begin to swell. For conditions and other formulas applicable to your locality consult your local County Agent.

ACCEPTED

8/8/71

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
OR ECONOMIC POISON REGISTERED
UNDER NO. 1388-23 SUBJECT
~~ATTACHED COMMENTS.~~

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT REUSE CONTAINER — DESTROY WHEN EMPTY