

VIII Outdoor

NUISANCE PESTS IN OUTSIDE AREAS—To control flies, ants, cockroaches, boxelder bugs, clover mites, spiders, ticks, chiggers and earwigs, use 8 fl. oz. in 3 gallons of water and spray thoroughly window frames, doors, sills, screens, porch and patio walls, ceilings, foundations, walks, garbage cans and cracks, under boards or other situations where insects may hide. Do not treat plants or animals with this dosage rate. Do not allow children in treated areas until surfaces are dry.

Use coarse drenching sprays for lawns, soils and other surface applications. Use medium fine mist sprays for trees and plants. Thorough coverage is important. Use a clean sprayer—flush with clean water after each use. Carefully measure and mix insecticide and water for maximum economy and effectiveness. Do not use on certain ferns, poinsetta, hibiscus, papaya and gardenia and orchids as injury may occur.

WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. For eyes get medical attention. Avoid contamination of feed and foodstuffs.

Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

Do not use or store near heat or open flame. Do not contaminate ornamental fish ponds.

FISH AND WILDLIFE—CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.



**DIAZINON® EMULSIFIABLE
Lawn and Garden Insecticide**

Contains Diazinon(R) Insecticide
Not for Use In The Home

INGREDIENTS

| | |
|---|------|
| ACTIVE INGREDIENTS: | |
| 0,0-Diethyl 0-(2-Isopropyl-6-Methyl-4-pyrimidinyl) phosphorothioate | 25% |
| Aromatic Petroleum Derivative Solvents | 57% |
| INERT INGREDIENTS: | 18% |
| Total | 100% |

(R) Trademark CIBA GEIGY CORPORATION
Ardsley, New York

WARNING

KEEP OUT OF REACH OF CHILDREN

Read Complete WARNING on Side Panel

EPA Reg. No. 1386-573

EPA Est. No. 904-NJ-1

NET CONTENTS
H&G-149

Distributed By

1 U.S. PINT
PROD. 2110

UNICO DIAZINON® EMULSIFIABLE

Contains 25.0% Diazinon(R) Insecticide
For Lawn and Garden Insect Control

Kills most outdoor insects which destroy lawns, roses, flowers, trees, shrubs, or annoy pets and people. Provides positive control of many "hard-to-kill" sucking and chewing insects. Works fast and remains effective for an extended period of time.

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF—To control chinch bugs, sod webworm, Bermuda mites, ants, ticks, earwigs and lawn bill bugs use 2 tablespoons (1 fl. oz.) in 3 gallons of water. Covers 125 square feet. Thoroughly wet down grass a few hours before applying. For lawn bill bugs, water lawn thoroughly after treatment.

ROSES AND FLOWERS—To control aphids, mites, leafhoppers and thrips on roses, azaleas, chrysanthemum, gladioli and carnations use 2 tablespoons (1 fl. oz.) per 3 gallons of water and spray plants till wet and dripping. Be sure to spray under sides of leaves.

TREES AND SHRUBS—To control aphids, mites, scale crawlers, leaf miners and bagworm on oak, maple, locust, holly, boxwood, pine, spruce and willow use 2 tablespoons (1 fl. oz.) per 3 gallons of water and spray to wet all leaves and branches to dripping point.

FRUITS—To control codling moths, mites, aphids, leaf rollers, scale crawlers on apple, peaches, oranges, plums, grapes, cherries, and strawberries use 2 tablespoons (1 fl. oz.) per 3 gallons of water. Wet the leaves and branches. Start spraying after blooming and repeat at 14-day intervals as needed. Do not spray fruit within 21 days of picking.

VEGETABLES—Controls Mexican bean beetles, aphids, leaf miners, cabbage worms, cucumber beetles on tomatoes, carrots, potatoes, beans, cucumber, onions, cantaloupes, peppers and cabbage. Use 2 tablespoons (1 fl. oz.) per 3 gallons of water. Will cover about 1,000 sq. ft. Applications are usually needed every 7 to 10 days. Do not apply within 1 days of picking tomatoes, 35 days of harvesting potatoes or 14 days of picking other crops listed.

To control cutworms, wireworms, root maggots, garden symphylans and mole crickets in carrots, cabbage, corn, lettuce, beans, pole beans, radishes, red beets, snap beans, tomatoes and turnips by soil treatment prior to planting, apply 1/2 fl. oz. per 100 sq. ft. in sufficient water to obtain thorough coverage. Then mix the spray thoroughly into the top 4 - 8 inches of soil.

UNIVERSAL COOPERATIVES, INC.
ALLIANCE, OHIO 44601