



APRICOTS, CHERRIES, FIGS, NECTARINES, PEACHES, PLUMS, PRUNES - Brown almond (Coville), European red, McDaniel, Pacific, peach silver, European, Schuler, two spotted, Willamette and yellow mites. Apply at petal fall to first cover or whenever mite population appears. Repeat application if necessary, wait 30 days to avoid excessive residues.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, OLALLIE BERRIES, RASPBERRIES Brown almond (Coville), European red, McDaniel, Pacific, two spotted, Willamette and yellow mites. Apply at petal fall or whenever mites appear and repeat if necessary. Use at the rate of 3 to 6 parts per acre. Do not apply within 2 days of harvest.

GRAPES, CHESTNUTS, FILBERTS, HICKORY NUTS, PECANS, WALNUTS - Brown almond (Coville), European red, McDaniel, Pacific, two spotted, Willamette and yellow mites. Apply at petal fall or whenever mites appear and repeat if necessary.

STRAWBERRIES Six spotted, strawberry (Atlantic) and two spotted mites. Apply when mites appear and repeat as necessary.

For cyclamen mites apply 2 to 3 thorough high pressure, drenching spray (400 gallons per acre) at 10 to 20 day intervals when new growth starts in spring or use as a spot treatment in infested areas during production season.

ADDITIONAL DIRECTIONS ON ATTACHED CARD

OTHER USES

CLOVER MITES IN BUILDINGS - Use 1 1/2 to 2 parts UNICO KELTHANE EMULSIFIABLE per 100 gallons. Spray walls, foundations, windowsills, plants and lawns at base of infested buildings.

LAWN AND TURF GRASSES Bermuda, brown and clover mites. Use 1 to 2 parts UNICO KELTHANE EMULSIFIABLE per 100 gallons (2 tablespoons per gallon). Apply 2 1/2 gallon dilute spray per 1,000 square feet. Make application when mites appear covering the entire lawn area with a thorough spray and repeat (10 to 14 days) if necessary.

CAUTION

Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mist. If inhaled or swallowed, if taken internally, get medical attention. Do not store in food or foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, ponds. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not use empty drums. Return to drum recycler or destroy by perforating, crushing and burying in a safe place.

NOTICE

Use only for the purposes and conditions specified on this label. Do not use for any other purpose.

**** KELTHANE EMULSIFIABLE**

ACTIVE INGREDIENTS:

*1,1-Bis(p-chlorophenyl)-2,2,2-trichloroethanol	18.5%
Xylene	73.0%
INERT INGREDIENTS:	8.5%
Total	100.0%

*Equivalent to 1.6 lbs. Active Ingredients Per Gallon
U.S. Patent #2,812,362 and 2,812,280

**Registered Trademark Rohm and Haas

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel
EPA Reg. No. 1386-575

NET VOLUME 1 U.S. PINT LIQUID
H&G-150E-2-73 DISTRIBUTED BY PROD. 2115

UNICO KELTHANE EMULSIFIABLE
DIRECTIONS FOR USE

UNICO KELTHANE EMULSIFIABLE gives high initial kill and long residual activity against most species of agricultural mites including the full width of the following:

Apple Rust	Peach Silver	Tomato Russet
Clover Mite (brown almond)	Plum Nursery	Topical
Cyclamen	Schöene	Two Spotted (Red) Spider
European Red	Two spotted	Willamette
McDaniel	Strawberry (Atlantic)	Yellow Carpini
Pacific		

UNICO KELTHANE EMULSIFIABLE is a specific miticide and will not kill bees and beneficial insect predators when used as directed.

UNICO KELTHANE EMULSIFIABLE can be applied alone or with other pesticides in recommended spray schedules. It should not be used with fungicides or with materials not compatible with solvent based pesticide formulations. For further details see specific recommendations on this label, then consult current State Agricultural Experiment Station spray schedules or State Extension Service Specialists for additional details on compatibility, dosage, timing and other factors.

UNICO KELTHANE EMULSIFIABLE is mixed with water and may be applied with conventional dilute or concentrate ground sprayers or with aircraft. Follow the dosage rates per acre listed below regardless of the gallonage of the spray applied. The higher dosages suggested are for larger trees or plants or heavier mite populations. Uniform coverage is necessary but the optimum spray coverage varies with the type of equipment, size and density of the crop plant. Use 5 to 10 gallons of spray per acre in aircraft.

To avoid excessive residue at harvest do not exceed the maximum dosage per acre or the time restriction between last application and harvest for each crop group listed in the table attached.

VEGETABLES AND FIELD CROPS

BEANS, CANTALOUPE, CUCUMBERS, HOPS, MELONS, PEPPERS, PUMPKINS, SQUASH (WINTER AND SUMMER), TOMATOES, AND WATERMELON - Pacific, Tomato Russet, Tropical and Two spotted mites. Apply at first signs of mite buildup and repeat as necessary. Do not feed crop residues to dairy or other animals. **DO NOT USE ON EGGPLANTS.**

FRUITS AND NUTS

APPLES, CRABAPPLES, PEARS, QUINCES - To clean up clover and European red mites and to control apple rust, McDaniel, Pacific, Schöene, two spotted, Willamette and yellow (Carpini) mites. Make application in second or third cover spray or whenever mite population appears. In areas where the preventive schedule is accepted practice, use two successive applications at 10-14 day intervals per 100 gallons dilute spray applied at 10-14 day intervals. Make thorough full coverage application. Note: UNICO KELTHANE EMULSIFIABLE may cause fruit spots on pears under certain conditions and should be used at the lower dosage rate with caution on this crop. To avoid possible injury do not use dilute formulations with UNICO KELTHANE EMULSIFIABLE in the cover sprays.

SUMMARY OF UNICO KELTHANE EMULSIFIABLE DOSAGE RATES IN PER ACRE AND LIMITATIONS BETWEEN LAST APPLICATION AND HARVEST

Crop	Dosage		Rate	
	Pts. Per Acre Min.	Pts. Per Acre Max.	Pts. Per 100 Gals. Dilute Sprays	Cutoff Days Before Harvest
Alfalfa and Clover (both grown for seed only use restricted to California, Idaho, Nevada, Oregon, and Washington)	3	6	3	
Chestnuts, Filberts, Hickory, Nuts, Pecans, Walnuts	8	20	2	14
Apples, Crabapples, Pears, Quinces	8	20	2	7
Apricots, Nectarines, Peaches	8	12		14
Beans (Dry)	3	4		45
Beans (dry green, lima)	2	3		7
Beans (dry green, lima) (California only)	4	5		21

UNITED CO OPERATIVES INC
ALLIANCE OHIO 44601

and lawns at base of infested buildings.

LAWN AND TURF GRASSES Bermuda, brown and clover mites. Use 1 1/2 to 2 pints UNICO KELTHANE EMULSIFIABLE per 100 gallons (2 tablespoons per gallon). Apply 2 1/2 gallon dilute spray per 1,000 square feet. Make application when mites appear covering the entire lawn area with a thorough spray and repeat in 10 to 14 days if necessary.

CAUTION

Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mist harmful if swallowed. If taken internally, get medical attention. Do not store near feed or foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, ponds. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not reuse empty drums. Return to drum conditioner or destroy by perforating, crushing and burying in a safe place.

NOTICE

Use only for the pests and conditions with which this product is registered. See warning statement on this label.

Total 100.00
 *Equivalent to 1.6 lbs. Active Ingredients Per Gallon
 U.S. Patent #2,812,362 and 2,812,280

**Registered Trademark Rohm and Haas

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1386-575

NET VOLUME 1 U.S. PINT LIQUID

H&G-150E-2-73 DISTRIBUTED BY PROD. 2115

VEGETABLES AND FRUITS
 BEANS, CANTALOUPE, CUCUMBERS, MELONS, PEPPERS, SQUASH, WATERMELONS
 SQUASH, WINTER AND SUMMER TOMATOES, AND THE FAMILY

UNITED CO OPERATIVES INC
 ALLIANCE OHIO 44601

SUMMARY OF UNICO KELTHANE EMULSIFIABLE DOSAGE RATES IN Pints PER ACRE AND LIMITATIONS BETWEEN LAST APPLICATION AND HARVEST

Crop	Dosage		Rate	Cutoff
	Unico Kelthane E. Pt. Per Acre Min.	Max.	Pts. Per 100 Gals. Dilute Sprays	
Alfalfa and Clover (both grown for seed only—use restricted to California, Idaho, Nevada, Oregon, and Washington)	3	6	3
Chestnuts, Filberts, Hickory, Nuts, Pecans, Walnuts	8	20	2	14
Apples, Crabapples, Pears, Quinces	8	20	2	7
Apricots, Nectarines, Peaches	8	12	14
Beans (Dry)	3	4	45
Beans (dry, green, lima)	2	3	7
Beans (dry, green, lima) (California only)....	4	5	21
Caneberries (all)	3	6	2	2
Cantaloupes, Cucumbers, Melons, Pumpkins, Squash, Watermelons	1 1/2	2	2
Cherries	8	12	2	7
Figs, Plums, Prunes	8	10	2	7
Grapes, Hops	3	6	2	7
Strawberries—Two-spotted and similar mites	2	4	4	2
Strawberries—Cyclamen mite	12	3	2
Tomatoes, Peppers	2	4	2
Clover Mites in Buildings	Full Coverage		1 1/2-2
Lawn and Turf Grasses	Full Coverage		1 1/2-2

*Gallonage of spray will vary with size and number of trees or plants per acre and the density of the foliage. In any case, thorough coverage of all parts of the plant is essential for good mite control.

Do not apply UNICO KELTHANE EMULSIFIABLE in any formulation on eggplant or avocados.

In general UNICO KELTHANE EMULSIFIABLE applications should be made as the various mite species begin to appear in threatening numbers and repeated as necessary for adequate seasonal control, except as noted or recommended locally.

ACCEPTED
 FEB 22, 1973
 FEDERAL WISE OF OHIO
 BY: [Signature]