



UNICO Fenthion Fly and Mosquito Spray

DIRECTIONS FOR USE

Read entire directions. Use strictly in accordance with label CAUTIONS, WARNINGS, AND DIRECTIONS.

UNICO FENTHION FLY AND MOSQUITO SPRAY is recommended for use wherever mosquitoes and flies are a problem, except in water where shrimp, crabs, or crayfish are of value. IT SHOULD NOT BE USED ON FOOD CROPS. DO NOT STORE IN OR AROUND THE HOME ENVIRONMENT.

It provides long residual control. The spray can be applied to most surfaces including freshly whitewashed area, without loss of biological effectiveness.

RESIDUAL SPRAY FOR FLY AND MOSQUITO CONTROL. For long residual control of mosquitoes and residual fly control, apply 4 to 8 fluid ounces in 1-gallon of water for application per 500 square feet or to run-off in localized area on patios, other outdoor structural surfaces, in buildings including dairy and beef barns and poultry houses. DO NOT APPLY DIRECTLY TO ANIMALS. Residual lasts 7 to 8 weeks, for the lower rate and even longer at the higher rate. Repeat when fly population appears to increase. DO NOT APPLY AS A SPACE SPRAY.

MOSQUITO CONTROL

Mosquito Larval Spray: Apply specified dosage, 3 fluid ounces (0.05 lbs. active) per acre in sufficient water or oil to obtain uniform coverage, or per 1/2 gallons of water or oil* for localized applications. Apply as a light uniform spray. Allow at least 3 weeks between applications.

Mosquito Thermal Fog: Mix 2 1/2 to 5 fluid ounces per gallon of oil*. Apply with standard fogging machines calibrated to deliver 40 gallons per hour at a machine speed of 5 m.p.h. to cover a swath width of up to 350 feet. Repeat as necessary.

*Kerosene type oils, fuel oils, diesel oils and other base oils suitable for insecticide use.

Mosquitoes Spray Mist: Apply 3 to 6 fluid ounces (0.05 to 0.1 lb. active) per acre. For mist-blower machines calibrated to deliver 100 gallons per hour traveling at a speed of 4 m.p.h. to cover a swath width of up to 350 feet, use 4 1/2 to 8 1/2 gallons per 100 gallon of water. Allow at least 3 weeks between applications.

PEST CONTROL ON TREES, SHRUBS AND FLOWERING PLANTS—FOR USE BY PEST CONTROL OPERATORS AND COMMERCIAL NURSERYMEN ONLY. To control ants, aphids, mites, leaf hoppers, armyworms, juniper scale, lecanium scale, and bagworms on the ornamental plants listed below:

Trees and Shrubs

Arborvitae
Ash
Aspen
Birch
Dogwood
Euonymus

Juniper
Maple
Oak
Spruce
Sweet Gum
Tulip Tree

Viburnum
Yew

Flowering Plants

Begonias
Chrysanthemums
Geraniums
Lilies
Marigolds
Peonies

Phlox
Petunias
Roses
Snapdragons
Verbena
Zinnias

DO NOT APPLY TO HAWTHORN, AMERICAN LINDEN, SUGAR MAPLE, OR THE ROSE VARIETY, DELIGHTFUL, BECAUSE OF POSSIBLE PLANT INJURY.

Mix 4 teaspoonfuls to 1 gallon of water (4 pints per 100 gallons). The resulting emulsion is suitable for use in any conventional hand-operated or power-operated spray equipment.

Spray when air is still to avoid drift. Do not treat plants in direct sunlight during the heat of the day or when the temperature is in excess of 90° F. Apply thorough coverage to upper and lower surfaces of leaves. Avoid wetting plants excessively—only a fine mist is required. Repeat as necessary.



FOR USE IN DAIRY AND BEEF BARN

CONTAINS BAYTEX®

ACTIVE INGREDIENTS:

Fenthion (0,0-Dimethyl 0-[4-(methylthio)-m-tolyl] phosphorothioate) 25.20%
Aromatic Petroleum Distillate 67.98%

INERT INGREDIENTS 6.82%

Total 100.00%

Baytex is a Reg. TM of the German Trademark proprietor, Leverkusen U.S. Patent No. 3,042,703

WARNING

KEEP OUT OF REACH OF CHILDREN

See Remainder of WARNING Statement on Side Panel

EPA Reg. No. 1386-584

Net Volume

5 U.S. Gallons Liquid

FC-1000B-5-73

PROD. 278

MANUFACTURED BY

UNIVERSAL COOPERATIVES, INC., ALLIANCE, OHIO 44601

