

UNICO LO-V 2,4,5-TP

WHERE TO USE:

UNICO LO-V, 2,4,5-TP WEED AND BRUSH KILLER is used on broad-leaved weeds in rice, turfgrass, and sugar cane, woody plants in rights-of-way, utility lines, fence rows, and similar non-crop areas, and on the weeds in drainage ditch banks, lakes, ponds, marshes, and other areas.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with eyes, skin or clothing. Flush eyes with clear water and get prompt medical attention.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT apply this product to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Minute quantities may cause severe injury by drift or vaporization. Avoid drift of spray and use a coarse spray which is less likely to drift. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

DO NOT store this product near fertilizers, seeds, insecticides or fungicides. DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose.

If stored below freezing, it may be necessary to warm to 40° F. and agitate before using. This does not effect the efficiency of the product.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

PLANTS CONTROLLED

Woody Growth:

Blackhack Oak
Ground Ivy
Mesquite
Pin Oak

Witch Hazel
Post Oak
Red Oak
Silt Cedar

Sand Shinnery Oak
Scrub Oak
White Oak
Wild Blackberry
Yucca

Aquatic Weeds:

Alligator weed
Arrowhead
Pickersweed

Rush
Spatterdock
Waterlily

Broad-Leaved Weeds:

Broomrape
Buckhorn
Chicory
Cocklebur
Common Chickweed
Crabgrass

Curled Dock
Curly Indigo
Dandelion
Lambsquarter
Witchgrass
Witchgrass

Plantain
Pokeberry
Ragweed
Sunflower
Wild Lettuce

UNICO LO-V 2,4,5-TP

DIRECTIONS FOR USE

Apply 2 quarts of product per acre to a 1/2 inch layer of water to be used in spraying. Mix well, then add 1/2 quart of diesel oil per 100 gallons of water. Shake container prior to mixing and agitate spray material before use.

Best results are obtained when weeds are actively growing. Repeat application if necessary.

WOODY GROWTH IN NON CROP AREAS

Ground Equipment: Mix 3 to 4 quarts of product with 1/2 quart of diesel oil per 100 gallons of water and spray woody growth up to 6 to 8 feet in height covering the ground surface. On growth up to 10 feet high, use 2 quarts per 100 gallons.

Airplane Application: For control of Post Oak, Blackberry, and other woody growth, add 2 quarts of product to 1 gallon of diesel oil and 1/2 quart of diesel oil per 100 gallons of water. Apply to 1/2 inch layer of water and apply to one acre.

Spot Treatment: Mix 2 quarts of product with 1/2 quart of diesel oil per 100 gallons of water and apply to one acre.

AQUATIC WEEDS

Use 2 quarts of product per acre with 1/2 quart of diesel oil per 100 gallons of water. Application should be made when leaves are fully developed above water. Do not apply to more than 1/3 to 1/2 of a lake or pond. Do not apply to streams, rivers, or other bodies of water where the oxygen content of water, killing fish.

WEEDS IN TURF

Mix 1/2 quart of product with 1/2 quart of diesel oil per 100 gallons of water and apply to one acre. Best kill is obtained with 1/2 quart of product per 100 gallons of water. Do not apply to newly seeded turf.

Do not use on newly seeded turf. Do not use on newly seeded turf. Do not use on newly seeded turf.

WEEDS IN RICE

Use 1/2 quart of product per acre with 1/2 quart of diesel oil per 100 gallons of water. Application should be made when rice is in the early stages of growth. It is advisable to delay spraying until the rice is in the early stages of growth. Do not apply to rice until the rice is in the early stages of growth.

SUGAR CANE IN LOUISIANA

For control of broadleaved weeds, mix 1/2 quart of product in 15 gallons of water for use in a row. Apply in the row approximately 1/2 inch before the seeds are expected to germinate. If cane is in the row, treat around daily following the application.

FLORIDA ONLY. For control of broadleaved weeds, mix 1/2 quart of product in 15 gallons of water and apply to one acre. Apply in the row when weeds are actively growing. Do not apply to the row until the weeds are actively growing. A second application may be required.

ACCEPTED

JUN 14 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 242-200 SUBJECT
TO ATTACHED COMMENTS.

UNICO®

MCPA 4 AMINE WEED KILLER

ACTIVE INGREDIENT:

Dimethylamine salt of 2-methyl-4-Chlorophenoxy acetic acid* 52.2%

INERT INGREDIENTS: 47.8%

TOTAL 100.0%

*Equivalent to 2-Methyl-4-Chlorophenoxyacetic Acid 42.6%.

*Equivalent to 4 Pounds of 2-methyl-4-Chlorophenoxy acetic acid per gallon.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional CAUTIONS

EPA EST. No. 1386-OH-1

EPA REG. NO. 1386-587

Net Volume

1 U.S. Gallon (3.785 Liters) Liquid

FC-100dA-12-78

PROD. 118

MANUFACTURED BY

UNIVERSAL COOPERATIVES, INC., ALLIANCE, OHIO 44601

CAUTION

This product may cause skin irritation. Harmful if swallowed. Avoid inhaling spray mist. Avoid contact with eyes, skin or clothing. In case of eye contact, flush eyes with clear water and get prompt medical attention. Clean contaminated clothing promptly after use. In case of skin contact, wash with soap and plenty of water.

WARNING

DO NOT apply this product to non-target and susceptible plants like beans and other legumes, cotton, flowers, grapes, ornamentals, vegetables and other plants by direct spray, drift or otherwise. Use coarse sprays as these are less likely to drift. Vapors from this product may injure susceptible plants in the immediate vicinity.

DO NOT contaminate water to be used for domestic or irrigation purposes.

DO NOT store this product near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate feed or feedstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose unless it has been cleaned and decontaminated carefully with a suitable chemical.

DO NOT reuse empty containers. Destroy by burying in an isolated non-crop area well away from potable water supplies.

DO NOT contaminate water by cleaning of equipment or disposal of wastes.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

UNICO MCPA 4 AMINE is a

UNICO MCPA 4 AMINE with MCPA.

Arrowhead	Lamb's
Cocklebur	Ear
Goatsbeard	Mint
Kochia	Mustard

Buttercup	Hemp
Canada Thistle	Hoary
Dandelion	Plantain

To insure uniform results spray evenly. Mix the recommended amount of gallons and for surface application as stated.

DO NOT MIX THIS PRODUCT with other herbicides. Best results are obtained when used under favorable conditions. Severe weather conditions affect economic or weed control results.

FLAX—Use 1/4 to 1/2 pint of product per acre where flax is 4 to 6 inches high. Higher rates may be required to control weeds.

RICE—Use 1 1/2 to 2 1/2 pints of product per acre. Do not apply to rice until approximately 65 days after seeding. Do not apply to rice under high (90° F and high humidity) conditions.

SMALL GRAINS (BARLEY, OATS, RYE, WHEAT)—Apply spray mix during early stages of growth. Do not apply to crops after the earing stage.

SMALL GRAINS (BARLEY, OATS, RYE, WHEAT)—Apply spray mix during early stages of growth. Do not apply to crops after the earing stage. Protective canopy of nurse crop may be present. Do not apply to small grains underseeded to a pasture.

GRASSES GROWN FOR SEED—Use higher rate where weed control is desired during the spring and early summer months.

ESTABLISHED GRASSLAND—Use higher rate for control of less desirable species. If desirable species are present, reduce rate to 1/2 pint per acre.

NOTE: If control is not satisfactory, reapply in some areas, bent, buffalo, etc.

NON-CROP AREAS (FENCES, DRIVEWAYS, etc.)—Apply 1/2 pint per acre in enough water to give target plants thoroughly during bloom. Do not apply after treatment or until chemical has dried.