

UNICO

DIRECTIONS FOR USE NOT FOR USE IN OR AROUND THE HOME

Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

BEANS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

BROCCOLI: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

CARROTS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

CELERY: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

COLLARDS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

CUCUMBERS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

EGGPLANTS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:


LETTUCE: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

MUSTARD GREENS: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

POTATOES: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

SPINACH: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:

SWEET CORN: Apply to plants in effective concentration. Coverage of plants should be uniform. Do not apply to plants which are stressed or which are in bloom. Observe the following directions for use:



ACTIVE INGREDIENTS:

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	32.5%
Heavy Aromatic Naptha	60.7%
INERT INGREDIENTS	6.8%
	100.0%

CONTAINS 3 POUNDS THIODAN PER GALLON.
*Thiodan U.S. Patent No. 2,799,685
Thiodan is a registered trademark of Canadian Hoechst Limited.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

See Side Panel for Additional WARNINGS.
EPA Reg. No. 1386-594
EPA Est. No. 1386-OH-1

DO NOT STORE AT TEMPERATURE BELOW 32° F.


Net Volume
FC-1007A-4-76

5 U.S. Gallons Liquid
PROD. 242

MANUFACTURED BY

UNIVERSAL COOPERATIVES, INC., ALLIANCE, OHIO 44601

SWEET CORN
planting
South Ce
imately 3
TOBA
qts. SEED
Larvae: U
field app
TOM
Aphid: 2
1-1/3 qts
gals. wat
ORN
100 gals.
"Bonnafor
on birch
SHAI
PINE
and again
SRU
May whe
BOX
when ins
and other
DOG
repeat in
TAXI
treatment
SUNF
at onset c
APPL
Rust Mite
to livesto
PEAC
Green Pe
acre. Less
qts. per
sprays to
PLUN
spray to
per acre.
CHER
not exce
in delay



NOTI
There is
WAR
in eyes,
eyes with
and befo
and load
the Minir
Institute
WAR
DO I
DO I
DO I
This
marshes
favor dri
Apply thi
This
Do n
CON
deep in r

Use

UNICO

THIODAN 3 EC

ACTIVE INGREDIENTS:

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	32.5%
Hevy Aromatic Naptha	60.7%
INERT INGREDIENTS	6.8%
	100.0%

CONTAINS 3 POUNDS THIODAN PER GALLON.

*Thiodan U.S. Patent No. 2,799,685

Thiodan is a registered trademark of Canadian Hoechst Limited.

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel for Additional WARNINGS.

EPA Reg. No. 1385-594

EPA Est. No. 1386-OH-1

DO NOT STORE AT TEMPERATURE BELOW 32° F.

Net Volume

FC-1007A-4-78

5 U.S. Gallons Liquid

PROD. 242

MANUFACTURED BY

UNIVERSAL COOPERATIVES, INC., ALLIANCE, OHIO 44601

SWEET POTATOES (0)—Sweet Potato Root Borer: 2-3 qt. per acre. Begin application shortly after planting or as soon as they first appear. Do not exceed 3 applications. In the Southeast and South Central States only: 1-1/3 to 2-3 qt. per acre. Work into the soil to a depth of approximately 3 inches making a single treatment. Do not apply to treated plants.

TOBACCO (5)—Fly Borer, Hornworm, Tobacco Root Borer: Apply 2-3 qt. per acre. Do not exceed 3 applications. SEED BED—Mix 2-3 qt. per 100 gal. of water and spray daily for 6-8 days. Larvae: Use 1-3 qt. per 100 gal. of water and spray part bed with 1 qt. spray per 100 gal. field application of 2-3 qt. per 100 gal.

TOMATOES (Field and Greenhouse)—Fly Borer, Colorado Potato Beetle, Fly Borer, Tobacco Root Borer: 2-3 qt. per 100 gal. of water. Hornworms: 2-3 to 1-1/3 qt. per 100 gal. of water. Tobacco Root Borer: 1-1/3 qt. per 100 gal. of water. Stink Bugs: 1 to 1-1/3 qt. per 100 gal. of water. Chrysanthemum Leafminer: 2-3 qt. per 100 gal. of water.

ORNAMENTALS (greenhouse and out of doors)—Aphid, Cyclamen Mite, Whitefly, Red Canker, Dogwood Borer: 100 gals. On chrysanthemums, leaf miners will be retained at airports before plants are shipped. Use "Bonnafon Deluxe", "Frid" and "Night" chrysanthemum as inquiry may be made. Do not apply on birch trees. In greenhouses, do not use on any insected milk or vegetable.

SHADE TREES (except Birch), SHRUBS—Aphid: 2-3 qt. per 100 gal. of water. Dogwood Borer: 1 qt. per 100 gal. of water. Apply in late April or early May and again if necessary in late fall. Do not use on any insected milk or vegetable.

PINES (Austrian, Jack, Red, Scotch, White)—Aphid: 2-3 qt. per 100 gal. of water. Dogwood Borer: 1 qt. per 100 gal. of water. Apply in late April or early May when aphids are present but before galls are formed.

SPRUCE—Spruce Gall Aphid (New England area only): 2-3 qt. per 100 gal. of water. Apply in late April or early May when aphids are present but before galls are formed.

BOX ELDER—Box Elder Bug: 2 qt. per 100 gal. of water. Dogwood Borer: 1 qt. per 100 gal. of water. Apply when insects first appear. Repeat as required. In the summer or fall spray tree trunks, limbs, and other places where bugs concentrate.

DOGWOOD—Dogwood Borer: 1 qt. per 100 gal. of water. Apply in early June and repeat in 10 to 14 days. Do not exceed 2 applications per year to the ground level.

LILAC—Lilac Borer: 2-3 to 1-1/3 qt. per 100 gal. of water. Apply in early June and repeat in 10 to 14 days. Do not exceed 2 applications per year to the ground level.

TAXUS—Taxus Bud Mite: 2-3 qt. per 100 gal. of water. Apply 3 to 5 applications beginning in May. Do not exceed 3 applications. Make first application at onset of bloom and make two more applications at 4 to 7 day intervals. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

SUNFLOWER (0)—Sunflower Moth: 1-1/3 qt. per acre. Do not exceed 3 applications. Make first application at onset of bloom and make two more applications at 4 to 7 day intervals. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

APPLES (30)—Apple Aphid, Pear Aphid: 2-3 qt. per 100 gal. of water. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

PEACHES (30), APRICOTS (30), NECTARINES (30)—Black Cherry Aphid, Black Cherry Aphid, Green Peach Aphid, Peach Twig Borer, Peach Tree Borer: 2-3 qt. per 100 gal. of water. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

PLUMS (7), PRUNES (7)—Lesser Peach Tree Borer: 1 qt. per 100 gal. of water. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

CHERRY (21)—Plum Root Borer: 2-3 qt. per 100 gal. of water. Do not exceed 3 applications. Do not apply to treated plants to livestock. Do not exceed 2 applications per year to the ground level. Do not apply within 50 feet of livestock.

ANTIDOTE

NOTE TO PHYSICIANS—Active ingredient is a central nervous system stimulant and may be used in treatment of depression. There is no specific antidote. Benzhexol is a sedative which may be used in treatment.

WARNING—Hazardous if swallowed. Avoid contact with skin and eyes. Do not breathe vapors. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and wash with plenty of water; for eyes, flush with water. Wash thoroughly with soap and water before eating or smoking; wear clean clothes. During commercial or professional operations, and before eating or smoking, wear clean clothes. During commercial or professional operations, wear a protective respirator.

WARNING—Workers entering treated areas within 24 hours of application should wear protective clothing. Do not store at temperatures below 32° F. Do not use in undiluted form. Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, snails, shells and other water life. Do not apply to streams, rivers, marshes and estuaries. Do not apply where it will be likely to kill. Do not apply where it will favor drift from areas treated. Do not apply to water by means of equipment. Do not apply to water.

WARNING—Workers entering treated areas within 24 hours of application should wear protective clothing. Do not store at temperatures below 32° F. Do not use in undiluted form. Do not use, pour, spill or store near heat or open flame.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings on this label.