

1386-606

PM 17

10/26/81

18

Pyrethrin

Multi-Purpose Insecticide

For Use On Livestock And In Homes, Institutions, Restaurants, Food Processing Plants, Dairies, And Outdoors For Mosquito Control.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Side Panel For Additional CAUTIONS

ACTIVE INGREDIENTS:


Pyrethrins	0.1%
Piperonyl Butoxide, Technical*	1.0%
INERT INGREDIENTS	98.9%
Total	100.0%

*Equivalent to 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds. DO NOT TRANSPORT OR STORE UNDER 32° F.

Net Contents: 1 Gallon

EPA Reg. No. 1386-006
EPA Est. No. 1386-0H-1

Not Reviewed. Registrant claims to be in accordance with the registration of Pesticide Product Guidance for Phase 2 Response. Page 2.12, Active to Inert Change in Status.



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply with Conventional Mechanical or Compressed Air Equipment

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites, and cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3 - 6 ounces per 1,000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the crawling insects, named above under SURFACE SPRAYING, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horn flies, houseflies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horseflies, and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

(Continued on Side Panel)

DIRECT

(Continued)

OUTDOOR APPLICATION

Adult Mosquito Control: Treat the entire area. For velocity does not exceed 5 vegetation where mosquito stagnant pools or ponds a gallons per acre (4 1/2 ounce Using truck-mounted com as Romist models 51C 91 a 300 foot swath, calibrate minute

STORAGE

Do not contaminate wa disposal. Do not store un

PESTICIDE DISPOSAL: F used according to label according to Federal, Sta Resource Conservation ar

CONTAINER DISPOSAL: dispose of in a sanitary lan local procedures

Use only for the purpo limitations, cautions or wa

902504-5-82