



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

1386-618

10/14  
file

23

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Nik Ramswick  
UNIVERSAL COOPERATIVES INC.  
P.O. BOX 460 7801 METRO PARKWAY  
MINNEAPOLIS, MN 55440

Subject: Label Amendment Submission of 07/28/93 in Response to PR Notice 93-7  
EPA Reg. No. 1386-618  
DIMETHOATE 267EC SYSTEMIC INSECTICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable  
Printed with Soy/Canola Ink on paper that  
contains at least 50% recycled fiber

2714

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (7505C)  
U.S. Environmental Protection Agency  
401 M Street, SW  
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Attachment

h10e

3714

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division

Nik Ramswick  
UNIVERSAL COOPERATIVES INC  
P.O. BOX 460 7801 METRO PARKWAY  
MINNEAPOLIS MN 55440

Comment for: EPA Reg Nr.1386-618  
DIMETHOATE 267EC SYSTEMIC INSECTICIDE

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

The Agricultural Use Requirements section must be located in a clearly separate box on the product labeling -- with lines or other graphic indicators (such as contrasting color) to separate the section from the surrounding text.

Remove the statement "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application." from its current position within the Agricultural Use Box and place it above the Agricultural Use Box.

The Non-Agricultural Use Requirements section must either be placed in a box or printed on the label in a contrasting color from the surrounding text.

The circled and asterisked statement(s) in the Non-Agricultural Use Requirements box does not belong. Delete the sentence(s) from the Non-Agricultural Use Requirements box and place it where it appeared in your original labeling. If there are no remaining requirements in the box after you delete the sentence(s), delete the entire Non-Agricultural Use Requirements box.

The REI in the amended label is much more restrictive than required. There may be a misunderstanding of the WPS guidelines for dimethoate, which allow a 48 hour REI for all crops.

4/2/14

= TEXT ADDED TO ACHIEVE  
WPS COMPLIANCE

DIMETHOATE 267 EC  
Systemic Insecticide

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
MAR 23 1994

Under the Federal Insecticide,  
Fungicide and Rodenticide Act  
as amended, for the pesticide  
registration. EPA Reg. No.  
1386-618

KEEP OUT OF REACH OF CHILDREN  
WARNING - AVISO  
Si usted no entiende la etiqueta, busque a alguien para  
que se la explique a usted en detalle. (If you do not  
understand the label, find someone to explain it to you  
in detail.)  
See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENT:	
Dimethoate [O,O-Dimethyl S-(N-methylcarbamoyl)	
methyl phosphorodithioate] .....	31.44%
INERT INGREDIENTS* .....	68.56%
Total .....	100.00%

Contains 2.67 Pounds Dimethoate Per Gallon. \*Contains Xylene.

Net Contents: 2 1/2 Gallons

EPA Reg. No. 1386-618  
EPA Est. No. 1386-OH-1



UNIVERSAL COOPERATIVES, INC., MINNEAPOLIS, MINNESOTA 55440

BEST AVAILABLE COPY

5 of 14

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

May be fatal if swallowed or inhaled. Harmful if absorbed. May cause irritation of eyes, nose, throat and skin. Do not get in eyes, on skin, or on clothing. Do not breathe vapor. Keep away from domestic animals and foodstuffs. Do not contaminate or apply directly onto feed or foodstuffs.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- \*Coveralls over long-sleeved shirt and long pants
- \*Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or viton
- \*Chemical-resistant footwear plus socks
- \*Protective eyewear
- \*Chemical-resistant headgear for overhead exposure
- \*Chemical-resistant apron when cleaning equipment, mixing, or loading
- \*For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)
- \*For exposures outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering controls statements:** When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

---

**USER SAFETY RECOMMENDATIONS**

- \*Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - \*Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 

**STATEMENT OF PRACTICAL TREATMENT  
ORGANOPHOSPHATE**

**If Swallowed:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious

Page 14

person. Get medical attention. If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists. If in Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

**NOTE TO PHYSICIAN:** This product may cause cholinesterase inhibition. Antidote is Atropine.

For Medical Emergency information, call 1-800-228-5635, extension 138.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Agricultural Extension services.

#### PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

---

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days.

7-8-14

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- \*Coveralls over long-sleeved shirt and long pants
- \*Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or viton
- \*Chemical-resistant footwear plus socks
- \*Protective eyewear
- \*Chemical-resistant headgear for overhead exposure

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Read the above reentry statement and the precautionary statements to workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information. "WARNING: Area treated with Dimethoate 267 EC (contains Dimethoate) on (date of application). Do not enter treated areas without appropriate protective clothing until sprays have dried. (Insert here Statement of Practical Treatment from label.)"

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

**APPLICATION EQUIPMENT** - Dimethoate 267 EC can be applied by any pressure sprayer that will give uniform distribution of the spray such as ground sprayers, aerial sprayers or portable sprayers.

**AERIAL APPLICATION** - Automatic flagging devices should be used whenever feasible. If human flaggers are employed they must wear the protective clothing and respirator specified on this label.

**GROUND APPLICATION** - Apply in sufficient water for good coverage unless otherwise stated. Avoid drift to adjacent crops.

**AIRPLANE APPLICATION** - Apply in 3 to 10 gallons of water per acre. Avoid drift to adjacent crops.

APPLICATION IN ORCHARDS - Do not graze livestock in treated orchards.

CROP	INSECT	RATE	DAYS BETWEEN LAST APPLICATION AND HARVEST
<b>FRUIT CROPS</b>			
Apples, Pears (bearing & non-bearing)	Aphids, Leafhoppers Pears Psylla Mites (except rust mites)	3/4 - 1 1/2 pints/100 gallons of water	Do not apply less than 28 days before harvest. Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom.  Do not graze livestock in orchards (groves) that have been treated.
Lemons Oranges Grapefruit Tangerines	Aphids	Ground Equipment: 3/4 to 1 1/2 pts. per 100 gals. water. Apply as an outside coverage spray. Aircraft Equipment: 3 to 6 pts. per acre in 15 to 20 gals. water.	Do not apply less than 15 days before harvest. Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom. Make no more than 2 applications to mature fruit. Do not use on citrus seedling. Do not enter treated groves within 4 days of last application.
	Mites (except rust mites)	Ground Equipment: 3/4 to 1 1/2 pts. per 100 gals. water. Apply as a thorough distribution coverage spray.	Do not graze livestock in orchards (groves) that have been treated.
	Scales (except black or snow)	Ground Equipment: 1 1/2 to 2 1/4 pts. per 100 gals. water. Apply as a thorough coverage spray.	Do not apply less than 45 days before harvest. Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom.  Make no more than 2 applications to mature fruit. Do not use on citrus seedling. Do not enter treated groves within 4 days of last application.  Do not graze livestock in orchards (groves) that have been treated.
	Thrips	Ground Equipment: 3/4 to 1 1/2 pts. per 100 gals. water. Apply as a mist spray. Aircraft Equipment: 3 to 6 pts. per acre in 15 to 20 gals. water.	Do not apply less than 15 days before harvest.  Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom.  Make no more than 2 applications to mature fruit. Do not use on citrus seedling. Do not enter treated groves within 4 days of last application.
	Whiteflies	Ground Equipment: 1 1/2 pts./100 gals. water. Apply as a thorough distribution coverage spray.	Do not graze livestock in orchards (groves) that have been treated.

BEST AVAILABLE COPY



<p>Cirrus (non-bearing) (Ariz. &amp; Calif.)</p>	<p>Thrips Aphids</p>	<p>FOLIAR SPRAY: Apply 1 1/2 pts. per 100 gallons of water. Repeat application as necessary. Do not enter treated groves within 4 days of last application. SOIL DRENCH: For trees 1-3 years old, apply 3 quarts per acre in the furrow or soil ring around the base of the tree. Apply as soon as insect injury to new growth begins. Use only on non-bearing trees. Nursery stock application should not be made in the year trees begin to bear fruit. Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom.</p>
--	--------------------------	--

**NUTS**

<p>Pecans</p>	<p>Aphids, mites, leafhoppers</p>	<p>Ground Equipment only: 1 pt./acre.</p>	<p>Do not apply less than 21 days before harvest. Do not graze livestock in treated groves.</p>
---------------	---------------------------------------	---	---

**VEGETABLE CROPS**

<p>Beans (green, lima, snap, dry)</p>	<p>Aphids, Leafhoppers, Leaf miners, Mites, Lygus bugs</p>	<p>3/4 - 1 1/2 pt./acre.</p>	<p>Beans may be harvested on day of application. Do not feed treated vines. This pesticide is highly toxic to bees. Do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom.</p>
<p>Broccoli Cauliflower</p>	<p>Aphids</p>	<p>3/4 - 1 1/2 pints/acre</p>	<p>Do not apply less than 7 days before harvest.</p>
<p>Cabbage</p>	<p>Aphids</p>	<p>3/4 - 1 1/2 pints/acre</p>	<p>Do not apply less than 7 days before harvest.</p>
<p>Head Lettuce</p>	<p>Aphids Leafhoppers Leaf miners</p>	<p>3/4 pint per acre</p>	<p>Do not apply less than 7 days before harvest.</p>
<p>Leaf Lettuce Spinach Collards Kale, Turnip (Greens &amp; roots), Mus- tard greens, Swiss Chard Endive (escarole)</p>	<p>Aphids Leafhoppers Leaf miners</p>	<p>3/4 pint per acre</p>	<p>Do not apply less than 14 days before harvest.</p>
<p>Melons (except Watermelons)</p>	<p>Aphids Leafhoppers Leaf miners Thrips</p>	<p>1 1/2 pints/acre</p>	<p>Do not apply less than 3 days before harvest.</p>
<p>Peas</p>	<p>Aphids</p>	<p>1/2 pint per acre</p>	<p>Peas may be harvested on day of application. Do not feed or graze hay within 21 days after last application when a stationary viner is used. Do not feed or graze when a mobile viner is used. This pesticide is highly toxic to bees. Do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom. Do not apply Dimethoate to peas more than once per season.</p>
<p>Peppers</p>	<p>Aphids Leaf miners Maggots</p>	<p>3/4 - 1 pint/acre</p>	<p>Peppers may be harvested on day of application</p>
<p>Potatoes Tomatoes Watermelons</p>	<p>Aphids Leaf miners Leaf hoppers</p>	<p>3/4 - 1 1/2 pints/acre</p>	<p>Do not apply less than 7 days before harvest</p>

**TEST AVAILABLE COPY**

Cyperus	Bactra moth larvae (apply as a drenching spray.)
Boxwood	Leaf miners, mealy bugs, mites.  For leaf miners, apply in spring, when leaf miner flies first appear or in early summer for control of larvae in the infested leaves.
Gardenias	Tea scale, whiteflies.
Gerberas	Thrips
Gladiolus	Aphids, thrips.
Poinsettia (Outdoor plants)	Mites, white flies, mealy bugs, aphids.
Ficus nitida (Outdoor plants)	Thrips (do not use on potted plants.)
Holly (English and American, not Burford variety)	Leaf miners, mites, soft scale.  For leaf miners, apply in spring when leaf miner flies first appear or in early summer for control of larvae in the infested leaves.

PLANTS	PESTS CONTROLLED
Hemlock	Mites, scale.
Roses (Outdoor plants)	Aphids, leafhoppers, mites, thrips.

Use 1/5 to 1/4 teaspoon in 1 quart of water to control the insects listed below.

Birch	Aphids, leaf miners.  For leaf miners, apply when leaves are expanded, about mid-May and repeat in early July.
-------	--

NOTE: 1 1/2 teaspoons per gallon is equivalent to 1/2 pint in 32 gallons of water.

Dimethoate 267 EC insecticide is effective generally in controlling aphids, thrips, leaf miners, scale insects, leafhoppers, and mites on certain ornamental plants. Apply sprays uniformly and thoroughly to foliage, except as otherwise stated, when insects first appear, or when damage is first observed, and repeat applications as needed. Do not overdose or overspray. Use only on the ornamental plants listed.

AVAILABLE COPY

11-8-14

FIELD CROPS			
Alfalfa (grown for hay)	Aphids Leafhoppers Lygus bugs Grasshoppers Reduction of Alfalfa Weevil Larvae	3/4 - 1 1/2 pts./acre	This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom. Do not apply within 10 days of harvest or pasturing. Make only one application per season. Effective only on cutting to which applied.
Alfalfa (grown for seed)	Aphids Leafhoppers Lygus bugs Grasshoppers Reduction of Alfalfa Weevil Larvae	3/4 - 1 1/4 pts./acre	This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom. Do not feed or graze livestock in treated crop hay, threshings or stubble within 10 days of application.
Cotton (grown in Ariz. & Calif.)	Lygus bugs Leafhoppers Thrips Black fleahoppers	3/4 - 1 1/2 pints/acre	Do not apply less than 14 days before harvest. Do not repeat applications within 14 days. Do not feed treated forage or graze livestock on treated fields. Do not apply more than 2 applications per season at the 1 1/2 pint rate.
Cotton (early season)	Aphids, Mites, Thrips, Fleahoppers	1/3 to 2/3 pint/acre	Do not apply less than 14 days before harvest. Do not repeat application within 14 days. Do not feed treated forage or graze livestock on treated fields.
	Lygus Bugs	2/3 pint/acre	
Cotton (Oklahoma, Texas)	Fleahoppers	1/3 to 2/3 pint/acre	Do not apply less than 14 days before harvest. Do not repeat application within 14 days. Do not feed treated forage or graze livestock on treated fields.
Field Corn	Banks Grass Mite (excluding Trans Pecos area of Texas)	1 - 1 1/2 pts./acre broadcast by air. Spray over the foliage when mites appear. Apply in ratio of 1 part DIMETHOATE 267 EC in 20 parts of water. Do not use less than 2 1/2 gals./acre for the 1 pt. rate and 3 3/4 gals./acre for the 1 1/2 pt. rate.	14 (foraging) 42 (grain) Apply as necessary. Make no more than three applications per year.  Do not apply to corn during the pollen-shed period.
Safflower (grown in Calif. & Ariz.)	Aphids Leafhoppers Lygus bugs Thrips	3/4 - 1 1/2 pints/acre	Do not apply less than 14 days before harvest. Repeat applications should not be made at intervals closer than 14 days. Make only 2 applications per season at the higher rate.

**BEST AVAILABLE COPY**

12-3-14

Sorghum (milo)	Aphids	<p>3/4 to 1 1/2 pts./acre GROUND APPLICATION: Apply above rates in 25 to 40 gallons of water per acre.</p> <p>AERIAL APPLICATION: Apply above rates in a ratio of 1 part Dimethoate 267 EC in 20 parts of water as follows: Rate of minimum gallons Dimethoate 267 EC water/acre.</p> <p>3/4 pt.                    2 1 pt.                      2 1/2 1 1/2 pt.                3 3/4</p> <p>Consult local extension agent for the proper timing of applications in your area.</p>	<p>Do not feed or graze within 28 days of last application.</p> <p>Make only 3 applica- tions per season as needed.</p> <p><b>DO NOT APPLY AFTER HEADING.</b></p>
-------------------	--------	--	---

Use 1 1/2 teaspoons in 1 gallon of water to control the insects on the ornamental plants listed below.

PLANTS	PESTS CONTROLLED
Azaleas (Outdoor plants)	Lace bugs, leaf miners, mites, tea scale, white flies.
Camellias	<p>Aphids, camellia scale, mites, tea scale.</p> <p>Foliar Spray: Apply 2 sprays, 6 weeks apart the first year, followed by annual applications soon after first growth begins in the spring.</p> <p>Soil Drench: Apply as a soil drench around the base of plants in early spring at the rate of 3 tablespoons in 1 gallon of water per plant up to 6 feet tall. Increase rate proportionately for larger plants.</p>
Carnations	<p>Aphids, thrips, mites.</p> <p>Soil Drench: Apply as a soil drench at the rate of 3 fluid ounces per 500 square feet of bed or bench (7 1/2 quarts per acre) in sufficient water for even distribution. Water in thoroughly after application.</p>

13 of 14

Use 3 teaspoons in 1 gallon of water to control the insects on the ornamental plants listed below.

PLANTS	PESTS CONTROLLED
Arborvitae	Aphids, bagworms, mites.
Cedar	Mites
Daylillies	Aphids, thrips.
Euonymus	Aphids, scale.
Iris	Aphids, iris borer, thrips.  For borer control, spray when new leaves are 5 to 6 inches tall.
Juniper	Aphids, bagworms, midges, mites.
Oak	Golden oak scale.
Pine	Aphids, bagworms, European pine shoot moth, Nantucket pine tip moth, Zimmerman pine moth.
Taxus (Upright or spreading yew)	Fletcher scale, mealy bugs, mites.

### CONTROL OF HOUSEFLIES

**RESIDUAL SPRAY** - To control houseflies in dairy barns, hog pens, calf barns, loafing sheds, poultry houses, and other farm outbuildings, apply a 1 percent residual spray to ceilings, walls, stanchions, and other surfaces on which flies congregate or rest. Prepare the spray by mixing 3 quarts in 25 gallons of water (1/2 pint in 2 gallons). Apply the spray to wet uniformly all fly-resting areas to the point of runoff. One gallon of spray will cover 500 to 1,000 square feet of wall or ceiling surface. Repeat applications as necessary.

Remove dairy animals, calves under one month of age, and poultry from buildings when applying residual sprays.

**SPOT SPRAY** - To control houseflies in localized areas, apply a spray made by mixing 5 fluid ounces (10 tablespoons) per gallon to surfaces around doorways and windows, in alleyways and feed rooms, and to other spots frequented by flies. Repeat as necessary.

**GOOD SANITATION IS A NECESSARY PART OF ANY EFFECTIVE FLY CONTROL PROGRAM.**

**MAGGOT SPRAY** - To control housefly maggots, mix 5 fluid ounces (10 tablespoons) per gallon and apply as a coarse spray or with a sprinkling can to fly-breeding media, such as poultry droppings in caged layer houses, garbage dumps and manure piles. Repeat applications if additional manure or garbage is added.

**DO NOT CONTAMINATE FEED OR FOODSTUFFS, DRINKING FOUNTAINS, FEED TROUGHS, AND LITTER. DO NOT USE IN MILK PROCESSING ROOMS, MILK HOUSES, AND MILK STORAGE ROOMS. DO NOT USE IN HOMES.**

**OUTSIDE SPRAY** - To control houseflies around home and recreation areas, animal quarters, food processing plants, warehouses, loading docks, and refuse areas, thoroughly spray exposed surfaces, such as walls, fences, garbage cans, and refuse containers with 1/2 pint in 2 gallons of water (2 tablespoons in 1 quart). Repeat as necessary.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Avoid temperature extremes. Shake well before using.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

CO-OP® is a Registered Trademark of Universal Cooperatives, Inc.  
Universal Cooperatives, Inc., Minneapolis, Minnesota 55440