



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 27 1994

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nik Ramswick
Universal Cooperatives, Inc.
P.O. Box 460
7801 Metro Parkway
Minneapolis, MN 55440

SUBJECT: Label Amendment Submission of 8/14/93 in Response to PR Notice 93-7
EPA Reg. No. 1386-626
Fruit Spray Powder

Dear Registrant:

EPA has determined that **this product does not fall within the scope of PR Notice 93-7**. The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.



Recycled/Recyclable
Printed with SoyCeres ink on paper that
contains at least 50% recycled fiber

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


James Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Enclosures

FRUIT SPRAY POWDER

A combination product for insect, disease, and mite control. One pound makes fourteen gallons of spray.

FOR HOMEOWNER USE ONLY

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 27 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

1386-626

KEEP OUT OF REACH OF CHILDREN

DANGER

See Back Panel For Additional Precautionary Statements
And Directions For Use

ACTIVE INGREDIENTS:

| | |
|-----------------------------------------------------------------------------------|---------|
| Malathion (0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate) | 7.50% |
| Captan* | 14.76% |
| Related Derivatives | 0.34% |
| Methoxychlor: 1,1,1-trichloro-2,2-bis (4-methoxyphenyl) ethane** | 18.25% |
| INERT INGREDIENTS | 59.15% |
| Total | 100.00% |

*N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide.

**Equivalent to 16.06% [1,1,1-trichloro-2,2-bis (p-methoxyphenyl) ethane] and 2.19% isomers and related products.

Net Contents: 1 Pound

EPA Reg. No. 1386-626
EPA Est. No. 1386-OH-1



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER**

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove contaminated clothing and separately launder clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush with plenty of water. Call a physician. **If Swallowed:** Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. **If Inhaled:** Remove person to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using, wear long pants and long sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing.

Apply as a spray by mixing with water. Use 7 level tablespoons per gallon of water. This rate applies to all crops listed on this label. Application rate per acre will depend upon tree or plant size, age, plant vigor and row width. For fruit trees, use 1 - 2 pints spray mixture on 5 - 8 foot trees; 1 - 2 quarts spray mixture on 8 - 14 foot trees; and 1 - 3 gallons of spray mixture on trees over 14 feet. The maximum permissible amount of Fruit Spray Powder which can be applied per acre is shown in parentheses "()" after crop listing.

Mix thoroughly before beginning to spray, maintain agitation while spraying, apply uniformly and thoroughly to foliage and fruits. Proper timing of applications and uniformly thorough coverage are essential. Avoid spraying when poor drying conditions are prevalent; foliage injury may result.

The following insects, mites, and plant diseases may be controlled by proper applications as indicated.

APPLES - Codling moth, apple maggot, plum curculio, redbanded leafroller, green apple aphid, rosy apple aphid, European red mite, leafhopper, two-spotted spider mites, Japanese beetle adults, scab, Brooks fruit spot, sooty blotch, black rot, bitter rot, tent caterpillars, pear slug, cankerworms, bud-moths, and crawler stages of San Jose oyster shell and scurfy scales.

Apply (1) just before bloom, (2) when most of the petals have fallen, (3) 7 days after petal fall, and (4) at 7 to 10 day intervals as needed in cover sprays. (27 pounds.)

The maximum number of foliar applications per growing season and/or maximum seasonal rate is 8 applications. Red Delicious and other sensitive apple varieties may be injured by early season applications.

PEACHES, APRICOTS - Oriental fruit moth, redbanded leafroller, slugs, European red mite, two-spotted spider mites, crawler stages of San Jose scurfy and oyster shell scales, brown rot, scab, leafspot, Japanese beetle adults, peach twig borer, and lesser peach tree borer.

Apply (1) when new buds show pink, (2) just before buds open, (3) when most of the petals have fallen, (4) 10 days after petal fall, and (5) on 7 to 10 day schedule, as needed. (Peaches 27 pounds; Apricots 16½ pounds.)

The maximum number of foliar applications per growing season and/or maximum seasonal rates are (1) Peaches - 8 applications at 2 lbs. ai/A/application, and (2) Apricots - 5 foliar applications using 2.5 lbs. ai/A/application.

STRAWBERRIES - Spittlebug, strawberry leafroller, leafhoppers, aphids, strawberry weevil, spider mites, Botrytis fruit rot, and leaf spot.

Apply (1) when growth starts in the spring, (2) at petal fall, and (3) at subsequent 7 to 10 day intervals as needed. (20½ pounds.)

The maximum number of foliar applications per growing season and/or maximum seasonal rate is 8 applications.

SPECIAL NOTES

Not effective if used with alkaline materials. Avoid spraying when poor drying conditions are prevalent. Some injury may occur on tender foliage.

STORAGE AND DISPOSAL

STORAGE: Store product in original container in a locked storage area. Do not store with or near feed or food products.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.