

# Formaldehyde Solution

**"DANGER"** Keep Out of Reach of Children  
See Back Panel For Additional Directions

**POISON** 

**WARNING:** Harmful if swallowed. Do not contaminate feed and foodstuffs. Keep away from children and domestic animals. Avoid contact

with skin. In case of skin contact, wash off with water at once. Avoid breathing vapor or fumes of Formaldehyde.

**ANTIDOTE:** Call a physician at once. When Formaldehyde is taken internally give large quantities of milk and raw eggs and induce vomiting. Keep patient quiet.


ACTIVE INGREDIENTS:

Formaldehyde .....	37%
Inert Ingredients - Water and methyl alcohol .....	63%
	100%

(Methyl alcohol less than 15%)

..... Liquid Measure

Packed by  
**KORINEK REMEDY COMPANY**  
Stayton, Oregon, U.S.A.



**THIS PRODUCT FOR  
AGRICULTURAL USE.**

U.S.D.A. Reg. No. 1388-9

## DIRECTIONS

For Grain Smut — mix one pint to forty gallons of water and dip partly filled sacks of grain until all is wet, or sprinkle over grain in piles and thoroughly mix until all is wet, then let drain and dry for twenty four hours before drilling.

For Potato Scab or Black Scurf — use one pint Formaldehyde to thirty-five gallons of water, soak potatoes in this solution for 1½ hours; do not cut them before soaking; let drain one hour before planting.

For conditions and other formulas applicable to your locality consult your local County Agent.